

₹10



# J I M A

Volume 64 (RNI) ♦ Number 01 ♦ January 2020 ♦ Kolkata

**JOURNAL** *of the* **INDIAN MEDICAL ASSOCIATION**

Official Publication of the Indian Medical Association

Indexed in

INDEX COPERNICUS  
INTERNATIONAL

Index Medicus

---

Volume 118 (JIMA) ♦ Number 01 ♦ January 2020 ♦ Kolkata

---



ISSN 0019-5847



Visit us at <http://www.ejima.in>

Newly Launched <https://onlinejima.com>

In patients with fast progression of **LUTS** in **BPH**

 **Veltam Plus**  
Tamsulosin MR 0.4 mg + Dutasteride 0.5 mg Tablets

The **smallest size** tablet to combAT **BPH**

In **BPH** for long term relief of **LUTS**

 **Veltam**  
Tablets  
Tamsulosin MR 0.2 mg & 0.4 mg

The **first tablet formulation** of **Tamsulosin** in India

In **Urinary Tract Infections**

 **Niftas**  
Nitrofurantoin 100 mg SR Tablets

The **best brand** of **Nitrofurantoin**

**INTAS PHARMACEUTICALS LTD.**

Corporate House, Near Sola Bridge, S.G. Highway, Thaltej,  
Ahmedabad-380054, Gujarat, INDIA • Website: [www.intaspharma.com](http://www.intaspharma.com)  
Email : [medical@intaspharma.com](mailto:medical@intaspharma.com)



**Goodness of Electrolytes**

**ENERZAL<sup>®</sup> ZERO**

**ZERO Calorie**

**Refreshing Taste**



## ADA\* Recommends<sup>1</sup>

Non-Nutritive sweeteners for cutting down the calorie intake

Sucralose has been extensively studied<sup>2</sup>

More than 110 safety studies reviewed & approved by FDA

Sucralose- accepted as safe by several national and international food safety regulatory bodies:

- US-FDA
- The Joint FAO/WHO Expert Committee Report on Food Additives.
- Food Standards Australia New Zealand.

1. <http://www.diabetes.co.uk/zero-drink.html>  
 2. <http://www.fda.gov/food/food-additives-ingredients/food-label-information-about-high-intensity-sweeteners-permitted-use-food-united-states>  
 \* American Diabetes Association

**Electrolyte Fluid without Calories**



**FDC Limited** 142-48, S.V. Road, Jogeshwari (W), Mumbai - 400 102

# ADMISSION NOTICE

Certificate & Diploma Under UGC Recognised University	UNDER WHO RECOGNISED FOREIGN UNIVERSITY	Eligibility
<ul style="list-style-type: none"> <li>▪ <b>Diabetology</b></li> <li>▪ <b>Ultrasound</b></li> <li>▪ <b>Rheumatology</b></li> <li>▪ <b>Radiology</b></li> <li>▪ <b>Pediatric</b></li> <li>▪ <b>Clinical Cardiology</b></li> <li>▪ <b>General Medicine</b></li> <li>▪ <b>Critical Care Medicine &amp; Many More.</b></li> </ul>	<ul style="list-style-type: none"> <li>☞ <b>MD / MS</b></li> <li>☞ <b>Master Of Medical Science</b></li> <li>☞ <b>MCH</b></li> <li>☞ <b>Diploma</b> ( In all traditional subjects)</li> </ul>	<h2>MBBS</h2>

## NATIONAL INSTITUTE OF MEDICAL SCIENCE

Trunk Road, Near Mawsumi Hospital & Research Centre  
Silchar -788001 Assam  
Affiliated By UGC & WHO recognized University  
For further details visit our website :- [www.nimssil.com](http://www.nimssil.com)  
E-mail : [drds20548@gmail.com](mailto:drds20548@gmail.com) / [contact@nimssil.com](mailto:contact@nimssil.com)  
Mobile -03842230152/09435072209/08811935789  
Admission forms are available on the website







## Stuck with just an MBBS?

Enroll in fellowship program with inbuilt Royal College (UK) training leading to Master of Medicine/ Master of Surgery



\*MTI placement assistance in UK

\*Scholarship available

\*Conditions Apply

**For More Details**

☎ 9108555231 | 📞 8056355149 | ✉ [antony.p@tauedu.org](mailto:antony.p@tauedu.org)  
[www.ucnedu.org/college-of-pg-medicine](http://www.ucnedu.org/college-of-pg-medicine)



## MBBS/MD/MS

### ARMENIA (EUROPE)

YEREVAN STATE MEDICAL UNIVERSITY  
(Founded in 1920)

- Leading Government Medical University
- International students from over 30 countries
- 1000 Indian students currently studying
- Regular Indian batches since 1987
- Wi-Fi campus with Robotics Lab & Simulation Centre
- State of Art Virtual 3-D Anatomy dissection table.
- 1100 Professors & 15 Clinical Hospitals
- Free Coaching for the 'Screening Test'
- Admission Test for U.G. in June / July
- Post Graduation in 27 Specialities
- 15 Clinical Fellowships (3-12 Months)
- 2 Sessions for P.G. Course : March & October

Annual Fee	First Year	2 <sup>nd</sup> Year & onward
Under-Graduate	5500 USD	5000 USD
Post-Graduate	6500 USD	6000 USD
Hostel Fee	700 USD/Year	

**Dr. Daljeet Singh Chauhan**  
 Official Representative of YSMU  
 President, YSMU Indian Alumni Association  
 D-136, 2<sup>nd</sup> Floor, Armenia Street,  
 Anand Niketan, New Delhi - 110021

📞 9810988669, 📠 (011) 2411-02-01




## www.ysmu.net



**Leaders in  
Diabetes &  
Cardiac  
Care in India**

**In T2DM,**

**Glycomet<sup>®</sup>-GP**

Metformin Hydrochloride 500/850/1000 mg SR + Glimepiride 0.5/1/2/3/4 mg

**Glycomet<sup>®</sup> S.R.**

Metformin Hydrochloride Sustained Release Tablets 500/850/1000 mg

**In the management of CAD,**

**Ecosprin<sup>®</sup> AV**

Enteric Coated Aspirin 75/150 mg + Atorvastatin 10/20 mg



There is nothing called 'away'  
**DISHA**  
is on your way

Launching  
**DISHA MAHANAGAR**  
at Rajarhat

[www.dishaeye.org](http://www.dishaeye.org)  
[dishaeyehospitals@gmail.com](mailto:dishaeyehospitals@gmail.com)



Barrackpore  
03366360000



Sinthi  
03366360005



Barasat  
03366360008



Berhampore  
03366360010



Durgapur  
03366360002



Behala  
03366360007



Mourigram  
03366360009



Burdwan  
03366360011



Teghoria  
03366360006



Gariahat  
03366360003



Sheoraphuli  
03366360004



Siliguri  
03366360001



Arambagh  
03366360014

**DISHA EYE HOSPITALS**

Largest Eye Care Provider  
in Eastern India

Appointment  
**033 6636 0000**



**Dr Rajan Sharma**  
National President,  
IMA



**Dr R V Asokan**  
Honorary Secretary  
General, IMA



**Prof (Dr) Jyotirmoy Pal**  
Honorary Editor,  
JIMA



**Dr Sanjoy Banerjee**  
Honorary Secretary,  
JIMA

# **JOURNAL** *of the* **INDIAN MEDICAL ASSOCIATION**

**Volume 118 (JIMA) ♦ Number 01 ♦ January 2020 ♦ Kolkata**

**ISSN 0019-5847**

## **CONTENTS**

### **Editorial :**

- ◆ End TB by 2025 : Difficult Mission, but Not Impossible  
— *Surya Kant* .....8

### **Original Articles :**

- ◆ A Retrospective descriptive study of snake bites in the medicine wards of Alipurduar hospital, West Bengal, India, from April 2013 to March 2014  
— *Hari Shankar Pathak, Tanuka Mandal, Tanushree Mondal, Partha Pratim Das, Uddalak Chakraborty, Tarun Kumar Paria, Purbasha Biswas* .....11
- ◆ A comparative study of Internal Jugular Vein Cannulation : Ultrasound Guided technique *versus* Conventional Blind technique — *Vaibhavi B Patel, Smita R Engineer, Kiran B Patel, Vikram P Mehta, Mona Bilandani* .....17

- ◆ Prevalence of Autoantibodies in patient complaining of multiple joint pain in a tertiary care hospital  
— *Sangeeta Das Ghosh, Mohiruddin Sk, Sandip Ghosh, Indrani Bhattacharyya* .....22

- ◆ Cross sectional analysis of suicidal behavior in adolescents — comparison with adult attempters  
— *P N Suresh Kumar, P K Anish* .....25
- ◆ Prevalence of anaemia in school going adolescents in a municipality town of West Bengal  
— *Ritesh Singh, Banasree Bhadra, Soumyajit Moulik, Suman Kumar Roy* .....30

### **Case Report :**

- ◆ Idiopathic Pancreatic Pseudocyst in a Child : a rare entity  
— *Shamita Chatterjee, Rajib Datta* .....35

# JOURNAL OF THE INDIAN MEDICAL ASSOCIATION

Founder Hony. Editor	: Sir Nilratan Sircar
Founder Hony. Business Manager	: Dr. Aghore Nath Ghosh
Hony. Editor	: Prof. (Dr.) Jyotirmoy Pal
Hony. Associate Editors	: Dr. Sibabrata Banerjee Dr. Sujoy Ghosh
Hony. Secretary	: Dr. Sanjoy Banerjee
Hony Assistant Secretary	: Dr. Shilpa Basu Roy
Ex-officio Members	: Dr. Iskandar Hossain, <i>Hony. Jt Finance Secretary, IMA (Hqs), Kolkata</i> Dr. Pijush Kanti Roy, <i>Hony. Joint Secretary, IMA(Hqs), Kolkata</i>

## OFFICE BEARERS OF IMA (HQS)

<i>National President</i>	<i>Honorary Executive Secretary</i>	<i>IMA Women Doctors Wing</i>
Dr. Rajan Sharma (Haryana)	Dr. Ajay Kumar (Bihar)	<i>Chairperson</i>
<i>Honorary Secretary General</i>	<b>JIMA (Calcutta)</b>	Dr. Mona P. Desai (Gujarat)
Dr. R. V. Asokan (Kerala)	<i>Honorary Editor</i>	<i>Honorary Secretary</i>
<i>Immediate Past National President</i>	Prof. (Dr.) Jyotirmoy Pal (Bengal)	Dr. Neeta S. Biyani(Maharashtra)
Dr. Santanu Sen (Bengal)	<i>Honorary Secretary</i>	<b>IMA Mission Pink Health</b>
<i>National Vice-Presidents</i>	Dr. Sanjoy Banerjee (Bengal)	<i>Chairperson</i>
Dr. Dipak Dhar Choudhury	<b>Indian Medical Journal (Delhi)</b>	Dr. Nilam Lekhi (Delhi)
(Uttarakhand)	<i>Chairman</i>	<i>Honorary Secretary</i>
Dr. Atul Durgeshankar Pandya	Dr. Ravi S. Wankhedkar	Dr. Vibha Tandon (Delhi)
(Gujarat)	(Maharashtra)	<b>IMA Junior Doctors' Network</b>
Dr. T. Narasinga Reddy (Telangana)	<i>Editor-in-Chief</i>	<i>Chairman</i>
Dr. G. N. Prabhakara (Karnataka)	Dr. Vedprakash Mishra (Maharashtra)	Dr. Adit Ketan Desai (Gujarat)
<i>Honorary Finance Secretary</i>	<b>Your Health (Calcutta)</b>	<i>Honorary Secretary</i>
Dr. Ramesh Kumar Datta (Delhi)	<i>Honorary Editor</i>	Dr. K. M. Abul Hasan (Tamil Nadu)
<i>Honorary Joint Secretaries</i>	Dr. Nandita Chakrabarti (Bengal)	<b>IMA Medical Students' Network</b>
Dr. Vijay Kumar Malhotra (Delhi)	<i>Honorary Secretary</i>	<i>Chairman</i>
Dr. V. K. Arora (Delhi)	Dr. Kakali Sen (Bengal)	Dr. Sreejith N. Kumar(Kerala)
Dr. Amrit Pal Singh (Delhi)	<b>IMA N.S.S.S. (Ahmedabad)</b>	<i>Honorary Secretary</i>
Dr. Pijush Kanti Roy (Bengal)	<i>Chairman</i>	Dr. Ajoy Kumar Saha (Maharashtra)
<i>Honorary Assistant Secretaries</i>	Dr. Kirti M. Patel (Gujarat)	<b>IMA National Health Scheme</b>
Dr. Usha Sridhar (Delhi)	<i>Honorary Secretary</i>	<i>Chairman</i>
Dr. S. K. Poddar (Delhi)	Dr. Yogendra S. Modi (Gujarat)	Dr. Ashok S. Adhao (Maharashtra)
<i>Honorary Joint Finance Secretaries</i>	<b>IMA N.P.P. Scheme</b>	<i>Honorary Secretary</i>
Dr. Dinesh Sahai (Delhi)	<i>(Thiruvananthapuram)</i>	Dr. Alex Franklin (Kerala)
Dr. Iskandar Hossain (Bengal)	<i>Chairman</i>	<b>IMA National Pension Scheme</b>
<b>IMA CGP (Chennai)</b>	Dr. Krishna M. Parate (Maharashtra)	<i>Chairman</i>
<i>Dean of Studies</i>	<i>Honorary Secretary</i>	Dr. Prashant H. Nikhade
Dr. Hiranmay Adhikary (Assam)	Dr. Jayakrishnan A. V. (Kerala)	(Maharashtra)
<i>Honorary Secretary</i>	<b>Apka Swasthya (Varanasi)</b>	<i>Honorary Secretary</i>
Dr. L. Yesodha (Tamil Nadu)	<i>Honorary Editor</i>	Dr. P. Gopeenathan (Kerala)
<b>IMA AMS (Hyderabad)</b>	Dr. Manoj Kumar Srivastava (Uttar Pradesh)	<b>IMA National Family Welfare Scheme</b>
<i>Chairman</i>	<i>Honorary Secretary</i>	<i>Chairman</i>
Dr. M. S. Ashraf (Tamilnadu)	Dr. Ashok Rai (Uttar Pradesh)	Dr. K. Vijayakumar (Tamil Nadu)
<i>Honorary Secretary</i>	<b>IMA Hospital Board of India</b>	<i>Honorary Secretary</i>
Dr. Mohan Gupta (Telangana)	<i>Chairman</i>	Dr. V. Sasidharan Pillai (Kerala)
<b>IMA AKN Sinha Institute (Patna)</b>	Dr. Vinod Kumar Monga (Delhi)	
<i>Director</i>	<i>Honorary Secretary</i>	
Dr. Yeshwant S. Deshpande	Dr. Jayesh M. Lele (Maharashtra)	
(Maharashtra)		



*Editorial***End TB by 2025 :  
Difficult Mission, but Not Impossible****Dr. Surya Kant**

*MBBS, MD (Gold Medalist), FCCP (USA),  
FAMS, FIAMS, FISEB, FNCCP, FCAI,  
FIMSA, FIAB, FICS, FUPDA, FIACM,  
FICP, FCGP*

*Professor and Head, Department of  
Respiratory Medicine, King George's  
Medical University, UP, Lucknow 226003*

*Whether we ourselves have escaped from the scourge of tuberculosis or not, there is probably hardly a family which has not had to do something with this dreaded disease.*

*— Pt. Jawahar Lal Nehru , 1st Prime Minister of India*

**T**uberculosis (TB), known since Vedic era in India as 'Kshaya Roga' or 'Raj Yakshma', finds similar description by Hippocrates as 'Pthisis' meaning to decay. On 24<sup>th</sup> March, 1882 that causative organism of TB was discovered by Robert Koch's, hence the disease is also known as 'Koch's disease' and 24<sup>th</sup> March is celebrated as 'World TB Day'<sup>1</sup>.

The basics of TB treatment are Nursing Care, Fresh Air and Good Nutrition, first described in Ayurveda and later on adopted by modern medicine as "Sanatoria Treatment". Chemotherapy era started in 1944, when Wakesmann discovered Streptomycin and since then few more drugs came into light, last being Rifampicin in 1980's. Then after a long gap of nearly 40 years Bedaquiline and Delamanid in recent years<sup>2-4</sup>.

Resurgence of TB occurred in late 80's, mainly due to Human Immunodeficiency Virus (HIV) and in 1993 World Health Organization (WHO) declared TB as 'GLOBAL EMERGENCY' due to rising problem globally, it's association with HIV and increasing drug resistance. Now it showed the incidences of TB as 140 per lakh population globally, whereas, it is 211 per lakh population in India. Deaths due to TB are 22 per lakh globally and 33 per lakh in India., HIV TB cases are 14 per lakh globally and 6.6 per lakh in India. The estimated Multiple Drug Resistance (MDR) or Rifampicin Resistant TB cases rested at 8.1 per lakh globally and in India it is 11 per lakh population<sup>5,6</sup>.

Approximately, half of the population is infected and one-fourth of the global annual new cases occur in India. Prevalence of TB is 3.2 million in India and incidences is 2.7 million new cases annually ie, more than 6000 people develop TB every day in India which amounts to more than 4 people developing TB every minute! About 4 lakh 23 thousands patients die due to TB every year which amounts to more than thousands deaths per day and one child everyday. There are about one lakh forty seven thousand patients of MDR-TB in India. Incidence of Multi Drug Resistant (MDR)-TB is 6.19% in all patients (2.84% in new and 11.6% in among previously treated)<sup>7,8</sup>.

If we look at the evolution of TB control in India, we find that National TB programme was first instituted in 1962. The programme was next reviewed in 1992 which saw that only 30% of patients were diagnosed and of these only 30% were treated successfully and hence the programme was declared as failed programme. These alarming figures gave birth to NTEP (Erstwhile RNTCP) pilot project in 1993. After the initial success, the NTEP project was scaled up in 1997 and finally on 24<sup>th</sup> March 2006, the entire country was covered under NTEP. The first National Strategic Plan for TB control was initiated for 2012-2017 to achieve universal access to quality TB diagnostics and treatment having guided activities and creating accountability against results. The Government of India, in May 2012, passed a mandate for all healthcare providers to notify every TB case diagnosed and/or treated, to local authorities. The national programme also rolled out an innovative and visionary electronic recording and reporting system (Nikshay) across the country in 2012. The need for mass-communication was addressed by the launch of Amitabh Bachchan's campaign, "TB Harega Desh Jeetega" on 24<sup>th</sup> March 2015. With this nationwide campaign of India Vs TB, our country has come a long way towards TB elimination<sup>9</sup>.

The real way forward has been shown to us by our Hon'ble Prime minister Shri Narendra Modi ji by organising the 'END TB SUMMIT' in Delhi on 13<sup>th</sup> March 2018 along with Dr Tedros Adhanom Ghebreyesus, Director-General of WHO.<sup>10</sup>

The Ministry of Health and Family welfare (MOHFW) in consultation with over 150 national and international experts working in the field of public health as well as private sectors finalized the new National Strategic Plan for TB 2017-2025 (NSP)<sup>11</sup>.

WHO definition of TB elimination implies reducing cases of TB to less than 1 per 10 Lacs population. Keeping this in mind, a vision of TB-Free India with zero deaths, disease and poverty due to TB is seen. The goal of this NSP is to achieve a rapid decline in burden of TB, morbidity and mortality while working towards elimination of TB in India by 2025. The requirements for moving towards TB elimination have been integrated into the four strategic pillars of "Detect – Treat – Prevent – Build" (DTPB).

Active case finding activity (ACF) among vulnerable groups is a focus over the next 5 years and considerable efforts are being made to reach these populations. The prioritization of vulnerable groups for screening and ACF has been formulated for urban, rural and tribal areas<sup>12-14</sup>.

Different choices for information and communication technologies (ICT) based treatment adherence support mechanisms like introduction of Real Time Medication Event Reminder Monitor (RT-MERM) and automatic pill counter have been introduced. The 'Nikhshay Poshan Yojana' launched by Government of India in March, 2018, aims to provide incentive of 500 rupees per month for the entire duration of treatment for nutritional support to TB patients. It has been implemented via a smart card linked to Aadhar which provides for a tamper proof storage of patient account and identity. To increase private sector involvement in TB control, every private doctor is being given incentive of Rs 1000 at the completion of treatment which he has notified.

Government of India declared TB a notifiable disease on 7th May 2012. Sticking to its earlier policy GOI took a harsher step towards TB control by issuing the gazette notification of MOHFW on 16<sup>th</sup> March, 2018 (published on 19<sup>th</sup> March). As per the gazette notification of MOHFW, clinical establishment – doctors, laboratories, pharmacists, chemists and druggists, that fails to notify TB patients will be booked under Sections 269 and 270 of the Indian Penal Code that carries a jail from six months to two years or a fine or both along with financial penalty. Government of India has taken all the steps after change of END TB by 2025 by our Honorable Prime Minister. However, malnutrition, Diabetes, Mellitus and Drug Resistance are the big challenges for this END TB mission<sup>15-17</sup>.

We, as a country have already eradicated polio. In the process we have learnt that having brand ambassador at state level, involvement of social, religious and local leaders in the awareness campaign, involvement of print and electronic media in spreading awareness, nukkadnataks, distribution of pamphlets, wall writings, folk dance etc to spread awareness in villages and remote places like tribal areas have brought us closer to our goal of eradication in much less time. We need to implement these learnings in our mission to eliminate TB also.

**The name of Revised National Tuberculosis Control Program (RNTCP) has been changed to National Tuberculosis Elimination Program (NTEP) on 30<sup>th</sup> December 2019. This Gesture is also in accordance with the Mission END TB 2025<sup>18</sup>.**

**We all should join hands to make India TB Free and**

**get rid of our country from this dreaded disease.**

**“END TB by 2025”.**

#### REFERENCES

- Koch R — Die Ätiologie der Tuberkulose, in: J. Schwalbe (Ed.), *Gesammelte Werke von Robert Koch*, Verlag von Georg Thieme, Leipzig, 1882; 1: 428-45.
- Schatz A — Streptomycin, a substance exhibiting antibiotic activity against Gram-positive and Gram-negative bacteria. *Proc Exp Biol Med* 1944; 55: 66-9.
- Sensi P — History of the development of rifampin. *Rev Infect Dis* 1983; 5: 3: 402-6.
- Patel RV, Riyaz SD, Park SW — Bedaquiline: a new hope to treat multi-drug resistant tuberculosis. *Curr Top Med Chem* 2014; 14 (16): 1866-74.
- WHO — TB A Global Emergency 1993. <https://apps.who.int/iris/bitstream/handle/10665/52639/WH-1993-Jul-Aug-eng.pdf?sequence=1&isAllowed=y>
- WHO — Multidrug and extensively drug-resistant TB (M/XDR-TB): 2010 global report on surveillance and response. Geneva: World Health Organization, 2010. [http://whqlibdoc.who.int/publications/2010/9789241599191\\_eng.pdf](http://whqlibdoc.who.int/publications/2010/9789241599191_eng.pdf) (accessed Aug 18, 2010).
- WHO — 2014-2016 National Anti-TB Drug Resistance Survey INDIA <https://tbcindia.gov.in/showfile.php?lid=3315>.
- Kant S, Maurya AK, Kushwaha RAS, Nag VL, Prasad R — Multi-drug resistant tuberculosis: An iatrogenic problem. *Bio Science Trends* 2010; 4(2): 39-47. 48-55.
- Nutritional Support to TB patients (NikhshayPoshan Yojana). Available from URL: <https://tbcindia.gov.in/showfile.php?lid=3318>. Last accessed 2018 on May 19.
- WHO Delhi TB Summit: WHO South-East Asia countries commit to intensified efforts, concrete progress to End TB <https://www.who.int/southeastasia/news/detail/13-03-2018-delhi-tb-summit-who-south-east-asia-countries-commit-to-intensified-efforts-concrete-progress-to-end-tb>.
- Revised National Tuberculosis Control Programme National Strategic Plan for Tuberculosis Elimination 2017–2025. <https://tbcindia.gov.in/WriteReadData/NSP%20Draft%202020.02.2017%201.pdf> 2017.
- WHO Active case finding: Systematic screening for active tuberculosis. <https://www.who.int/tb/areas-of-work/laboratory/active-case-finding/en/>
- Uplekar M, Cresswell J, Ottmani S, Weil D, Lönnroth K — Programmatic approaches to screening for active tuberculosis. *Int J Tuberc Lung Dis* 2013; 17: 1248-56.
- WHO Global Report on The End TB Strategy [https://www.who.int/tb/post2015\\_strategy/en/](https://www.who.int/tb/post2015_strategy/en/)
- Kant S, Gupta H, Ahluwalia SA — Significance of nutrition in pulmonary Tuberculosis. *J of critical reviews in food science and nutrition* 2015; 55: 955-63.
- Kant S, Hem Lata, et al. Verma Diabetes Mellitus with Pulmonary Tuberculosis A Double Trouble. *J Indian Med Assoc (JIMA)*, review article 2013; 111(3): 187-91.
- Narain A, Verma AK, Srivastava A, Kant S — Drug Resistance Pattern in the Clinical Isolates of Pulmonary Tuberculosis. *Acta Scientifica Microbiology* 2019; 2(10): 66-73.
- Government of India Ministry of Health and Family Welfare Nirman Bhavan, New Delhi, D.O. No. A. 66013/34/30.12.2019-TB.

#### Disclaimer

The information and opinions presented in the Journal reflect the views of the authors and not of the Journal or its Editorial Board or the Publisher. Publication does not constitute endorsement by the journal.

JIMA assumes no responsibility for the authenticity or reliability of any product, equipment, gadget or any claim by medical establishments/institutions/manufacturers or any training programme in the form of advertisements appearing in JIMA and also does not endorse or give any guarantee to such products or training programme or promote any such thing or claims made so after.

— *Hony Editor*

## Original Article

## A Retrospective descriptive study of snake bites in the medicine wards of Alipurduar hospital, West Bengal, India, from April 2013 to March 2014

Hari Shankar Pathak<sup>1</sup>, Tanuka Mandal<sup>2</sup>, Tanushree Mondal<sup>3</sup>, Partha Pratim Das<sup>4</sup>, Uddalak Chakraborty<sup>5</sup>, Tarun Kumar Paria<sup>5</sup>, Purbasha Biswas<sup>5</sup>

Snake bite is an important cause of accidental death in modern India and its public health importance has been underestimated. West Bengal is one of the high snake bite prevalence states of India. Alipurduar district, a heavily riverine forested district with huge areas of agricultural land, large number of tea gardens reports a considerable number of snake bites annually and 13 species of venomous snakes have been reported from here. No data on snakebite epidemiology, clinical profile and mortality in Alipurduar district is available yet. We conducted a hospital based retrospective descriptive study of 160 patients admitted with history of snake bite in the medical wards of the Alipurduar District Hospital over a period of 12 months. Details of the history including time and place of bite, clinical findings including local and systemic features of envenomation, tests performed, provision of Anti-Snake Venom and supportive medications and outcomes were obtained from the hospital records. 75% of the patients were from rural areas including tea gardens. 71.2% were males and 70% of the persons in the 30-49 years age group were affected. Seasonal variation in snake bite was seen with a peak in June-July. 70% of the snakebites had no features of envenomation and 60% of the bites occurred during the day time the lower limbs being bitten the most (83%). 70% of the snake bite cases were hospitalized within 6 hours of the bite and 88% of them attended government hospital without visiting any traditional healer. 80% of the envenomed cases had neurotoxic features. More than 70% cases had panic attacks and 80% had tight ligature with complications. Community education on prevention of snake bites, appropriate first aid following snake bites and appropriate cum adequate training of medical personnel in syndromic management of snakebites could reduce snake bite morbidity.

[J Indian Med Assoc 2020; 118(1): 11-6]

**Key words :** Snake bite, Types of envenomation, Clinical findings of snake bite.

Snake bite was included in the list of neglected tropical diseases by the World Health Organisation in the year 2009, removed in 2013 and then again reincluded in 2017<sup>1,2</sup>. 'Million Death Study' conducted by the Registrar General of India (RGI) estimates around 50,000 snakebite deaths per year in India based on "routine, representative re-

sampled household interview of mortality with medical evaluation (RHIME)"<sup>3</sup>. Snake bite is an important cause of accidental death in modern India and its public health importance has been underestimated with the morbidity estimated to be more than 30 – fold higher than the number declared from official hospital returns<sup>3,4</sup>. West Bengal is one of the high snake bite prevalence states of India besides Andhra Pradesh, Kerala, Tamilnadu and Maharashtra<sup>5</sup>. Incidence of snake bite depends on the frequency of contact between snakes and humans; snake bite occurs when humans move to the habitat of snakes like paddy fields, tea and coffee plantations or when snakes frequent domestic and peri-domestic areas in search of preys like rodents, fowls, etc or during flooding<sup>6</sup>. Alipurduar district which encounters large number of snake bites averaging 600-700 annually, is a heavily riverine forested district with famous wild life sanctuaries and has huge areas of agricultural land and a large number of tea gardens, all of

Department of General Medicine, College of Medicine and JNM Hospital, Kalyani 741235

<sup>1</sup>MBBS, MD (Gen Med), FICP, Professor

<sup>2</sup>MBBS, MD (Community Medicine), Senior Resident, Department of General Medicine, R G Kar Medical College and Hospital, Kolkata 700004

<sup>3</sup>MBBS, MD (Gen Med), MACP, Assistant Director of Medical Education, Government of West Bengal & Associate Professor, Medical College and Hospital, Kolkata 700073 and Corresponding Author

<sup>4</sup>MBBS, MD (Gen Med), FICP, Consultant Physician, Alipurduar District Hospital, Alipurduar 736121

<sup>5</sup>MBBS, Postgraduate Student of MD, Department of General Medicine, R G Kar Medical College and Hospital, Kolkata 700004

which are home to many species of snakes<sup>7,8</sup>. 25 species of snakes are reported to be common in undivided Jalpaiguri district, including Alipurduar (a separate district since June 2014) of which 13 are venomous and 12 non venomous, including Pythons<sup>7</sup>. The common venomous found during an ophiofaunal survey are Naja (Spectacled cobra), Bungarus caruleus (Krait), Naja Hannah (King cobra) and Trimeresurus species (Pit viper)<sup>7</sup>. No hospital based or community based surveys on snake bite epidemiology, clinical profile and mortality in Alipurduar District in the Northern Part of West Bengal, India is available yet. Deaths due to snake bite in India are mainly due to lack of awareness in the community on proper first aid, delay in reaching the hospital, where AVS is available, killing valuable time at the traditional healer's (Ojha) place, lack of confidence of doctors in using AVS infusion, improper protocol for its management, less stress on syndromic approach by doctors and sometimes lack of infrastructure for critically ill patients<sup>9,10</sup>.

#### AIMS AND OBJECTIVES

We aimed to study the epidemiology and clinical aspects of snake bites in and around Alipurduar, West Bengal to identify frequency at the district hospital, which is best equipped to manage snake bites; types of envenomation; sites of bites; signs and symptoms of envenomation; first aid provided at the community level and responses to provision of treatment as per the National and State Snake Bite Management Protocol at the district hospital and suggest measures to strengthen snake bite management in the district.

#### METHODS AND MATERIALS

We conducted a hospital based retrospective descriptive study of 160 patients admitted with history of snake bite in the medical wards of the Alipurduar District Hospital over a period of 12 months (April 2013 to March 2014). We recorded the history of snake bite including time and place of bite, species of snake involved, clinical findings including local and systemic features of envenomation, tests performed including the 20 minutes Whole Blood Clotting Test (20 WBCT), provision of Central Research Institute Polyvalent Anti-Snake Venom based on syndromic approach and Injection Atropine, Injection Neostigmine and Injection Adrenaline as outlined in the National and State Snake Bite Management Protocol as well as outcomes from the hospital records of Alipurduar District Hospital with the consent of the Superintendent of the hospital.

#### RESULTS

Of the total 160 cases admitted with history of snake bite in the medical wards of the Alipurduar District, 114 (71.2%) were males, 36 (28.8%) were females, 112 (70%) were in the age group of 30-49 years, 31 (19.4%) in between

50 to 70 years of age and 17 (10.6%) between 11 to 29 years of age (Fig 2). 120 (75%) of the patients were from the rural areas, mostly from tea gardens, agricultural and forest villages.

30 (18.8%), 24 (15%) and 10 (6.3%) were homemakers, farmers and forest workers respectively while the remaining 96 (60%) included people of various occupations, including 8 (5%) snake charmers and non-working persons.

90 (56%), 41 (25.6%), 19 (11.9%) and 10 (6.3%) of the cases occurred in the months of February-April, May-July, August-October and November respectively (Fig 3).

32 (20%) of the cases were brought within 3 hours of the snake bite, 80 (50%) brought between 3 and 6 hours of the bite and 48 (30%) cases brought after 6 hours of the snake (Fig 4).

96 (60%) of the snake bites took place during the daytime while on outdoor activities, including farming, playing, working in the forest and handling snakes while 32 (20%) snake bites took place at night, mostly on stepping on snakes unknowingly. 32 (20%) of the bites took place when the patients were asleep (Table 1).

134 (83.8%), 16 (10%) and 10 (6.3%) of the bites were in the lower limbs, upper limbs and other parts of the body respectively. 132 (82.5%) of the snake bite victims, including the snake charmers, could not identify the snakes (Fig 4). In 8 (5%) of the cases snakes were brought for identification and 3 cases were identified to be python bites.

128 (80%) of the snake bites cases were given tight ligature jeopardizing blood circulation, as well as, improper first aid 140 (88%) of the snakebite victims were brought straight to Alipurduar District Hospital or to other government hospitals and then referred to the Alipurduar District Hospital without contacting any traditional healer.

112 (70%) of the snakebite cases had no features of envenomation. Bite marks with or without fang (two puncture marks) marks were present in 112 (70%) of the cases. 83 (51.9%), 89 (55.6), 96 (60%) and 54 (33.8%) had local oedema, bleeding (generalized or from bite wound), pain and numbness respectively. 32 (20%) of the cases had cellulitis/compartamental syndrome/necrosis at site of bite due to tight ligature and injury and required referral for surgical intervention (Figs 1&5).

128 (80%) cases had panic attacks, including palpitation, chest pain, psychogenic dyspnoea, tremor, etc. 24 (15%) had neurological signs and symptoms, including ptosis, weakness of the limbs, aphasia, aphonia, etc. 5 (3.12%) had haematological signs and symptoms, including bleeding, ecchymosis. In 10 (6.3%) cases, 20 minute WBCT was positive. 1 (One) patient had renal complications with hematuria, hemoglobinuria, etc.

On an average 10 to 30 vials of AVS were used in 96

(60%) snake bite cases (without skin test) based on the syndromic management as per the government guidelines. 48 (30%) cases were administered Injections Atropine (0.6mg to 6mg) and Neostigmine (1.5mg to 9mg) at 30mins to 60mins interval. Patients were kept and observed for a minimum 48 hours in hospital. Anaphylactic shock was noted in 12 (7.5%) cases, all of whom responded to Injection Adrenaline. Minor reactions, including itching, rash, nausea, pain abdomen, vomiting were seen in 64 (40%) of the cases who received Injection AVS. No death occurred due to AVS reaction. 144 (90%) of the snakebite cases were discharged as cured and healthy. 12 (7.5%) were referred to a higher center while 2.5%) patients died at Alipurduar District Hospital. The discharged patients were all counselled regarding snake bite prevention, proper first aid and encouraged for early access to health care intervention at government hospitals, including Rural Hospitals & Primary Health Centres.

**DISCUSSION**

There was high number (70%) of dry bites or bite by non-poisonous snakes. Various studies suggest that envenomed bites constitute between 12% and 50% of the total number of bites and bites by non-poisonous snakes might be as high as 70% of all snake bites<sup>9-11</sup>.


Males were found to be more affected (above 70%) and that more than 70% of middle aged persons with age ranging from 30 to 49 years were affected. In Bankura District of West Bengal more than 50% of the snake bite victims were in the 30 – 50 years aged group<sup>12</sup>. In Paschim Midnapore District of West Bengal too the majority age group (>42%) affected was the 31 -50 age group<sup>13</sup>. Similarly, males accounted for more than 56% in Paschim Midnapore<sup>13</sup>. This may be explained by the fact that males and people in the 30-49 age group commonly remain engaged in outdoor activities in farms, tea gardens and forests<sup>6</sup>.

There was obvious seasonal variation with rising number of snake bites in the rainy season from May and

with highest incidence in July-August and decrease in December to pre-monsoon level<sup>3,6,10,12</sup>. Snake bite and death rate is always high during the rainy season<sup>12,14</sup>. Increased snake bites during the rainy season are probably due to flooding of the natural habitats of the rats and the snakes and their increased presence in and around human residences<sup>6,10,14</sup>. This probably explains the high number of snakebites among homemakers in Alipurduar. Bites during agricultural/tea plucking activities, which increase during the rainy season is probably another reason for high number of snake bites and deaths during this season<sup>10</sup>.

Bites occurred mostly during outdoor activities (60%) but bites during sleep were remarkably low at 20%. In a hospital based study at Bankura District of West Bengal more than 71% of the snake bites occurred during outdoor activities<sup>6,12</sup>. Common Krait bite is not very common in Alipurduar District though 4 (four) cases were suspected to be Krait bites based on the history of bite during sleep and mysterious clinical features of abdominal pain, sore throat and fever in the absence of bite marks and administered AVS to which they responded favorably.

Bites affected lower limbs (83.75%) predominantly. In tropical countries lower limb is affected more because of occupation, lack of use of protective footwear or less visibility in rural area due to lack of light<sup>6,12-14</sup>.



Before Treatment



After Treatment



Fig 1 — Necrosis at the site of snake bite

Table 1 — Time and Site of Snakebites, 2013-14, Alipurduar District Hospital, Alipurduar, West Bengal, India	
Time of Bite	Number of bites
Day time	96
Night time	32
During sleep	32
Area of bite	Number of bites
Lower limb	134
Upper limb	16
Other areas	10




Fig 2 —Age distribution of Snakebites, Alipurduar District Hospital, Alipurduar, West Bengal



Fig 3 — Monthly distribution of Snakebites in 2013-2014, Alipurduar District Hospital, West Bengal

More than 80% of the cases presented with panic attacks/early psychiatric features. Similar hospital based study of 121 snake bite victims in Bangladesh found that more than 80% of the patients had early psychiatric features, including severe anxiety, hallucination, irritability, psychogenic convulsion, difficulty in concentration, aggressive behavior, etc<sup>15,16</sup>. As per cognitive behavior is concerned, the snake bite could be a critical incident acting on existing psychological conditions including myths, misconceptions and ophidiophobia, triggering negative automatic thoughts leading to anxiety and depression<sup>17</sup>. Anxious people may over-breathe so that they develop pins and pin and needles of extremities, stiffness or tetany of their hands and feet and dizziness, vasovagal shock and high blood pressure and pulse with consequent adverse effects<sup>17</sup>. Reassurance that death is not imminent and that medical care is available is a very important first aid intervention. This will drive away their fear and excitement, slow the patient's heart rate and reduce the spread




Fig 4 — Time gap between Snakebite and Hospitalisation, 2013-2014, Alipurduar District Hospital, West Bengal

of venom<sup>15</sup>.

Improper first aid especially tight ligatures (20%) resulted in complications including cellulitis and compartmental syndrome. Tourniquets and compression bandages are potentially dangerous as they can cause gangrene, increased fibrinolysis and bleeding in the occluded limb, peripheral nerve palsies and intensification of local envenomation<sup>18</sup>. Traditional tight (arterial) tourniquets if applied tightly around the upper part of the limb are extremely painful as the limb becomes ischaemic and are very dangerous if left in place for long periods resulting in gangrenous limbs. <sup>15</sup> This practice is also strongly discouraged and only immobilization is recommended<sup>15,18</sup>.

70% of the snakebite victims reached the hospital within 6 hours of the bite. In a hospital based study in Malaysia, less than 50% snakebite victims reached the hospital in 24 hours since snakebite<sup>19</sup>. In Purulia District in West Bengal, less than 30% snakebite victims reached the hospital within 6 hours of bite<sup>20</sup>. This reflects that the awareness level and service access in Alipurduar District is relatively better.

Only 12.5% of the patients sought interventions from




Fig 5 — Local freatures in Snakebite cases, 2013-2014, Alipurduar District Hospital, West Bengal

traditional healers. A retrospective analysis of snakebite data from a community based epidemiological survey in South 24 Parganas in West Bengal, India, showed that 62% of snakebite deaths were due to intervention of traditional healers<sup>14</sup>. Among the snake bite patients in Burdwan District in West Bengal State, India 65.47% went to the traditional healers (ojhas) and 8.46% persons went to hospital only after consulting the ojhas.<sup>5</sup> This reflects that the people in Alipurduar are conscious about early hospitalization.

Local features were more common than systemic manifestations and in many were progressive suggestive of poisonous snake bites. Neurotoxic snake bites (15%) were more compared to vasculotoxic snake bites (3.12%), which is very much identical with the findings in Purulia district of West Bengal<sup>19</sup>. In Bankura district of West Bengal around 50% of snakebite deaths were due to neurotoxic envenomation<sup>12</sup>. However, in Paschim Midnapore district of West Bengal vasculotoxic bites were more than 82%<sup>12</sup>. The very low % of vasculotoxic bites is corroborated by the fact that snakes of the viperidae family are rare in Jalpaiguri District, including Alipurduar<sup>7</sup>.

Though more than 70% of the cases admitted with a history of snakebite were found to have dry bites or had unknown bites, 60% had been administered Injection AVS. This was likely to be primarily due to combination of various factors including predominance of history of snakebite by the victims or their relatives, local symptoms simulating venomous snakebites and lack of confidence on the part of the medical officers in ascertaining the signs of envenomation, high proportion cases having panic attacks, as well as, the apprehension of public backlash just in case the condition of the patient worsened reflecting lack of proper training in snake bite management. Low level of recognition of signs of envenomation and undue importance to local symptoms leads to administration of AVS to non-envenomed victims<sup>9</sup>.

The average use of 10 to 30 vials of AVS was as per the existing snakebite management protocols and corroborated with studies in other parts of the state, country and the world<sup>21-23</sup>. Though there were some cases of anaphylactic shock which was managed satisfactorily with Injection Adrenaline, no complications or death occurred due to administration of Injection AVS.

Syndromic approach based on state snake bite management protocol was helpful in treating more than 90% of the cases successfully even in the absence of identification of kind of snake involved. A syndromic approach is very useful when snakes have not been identified and clinical features caused by venoms of different species may overlap.<sup>15</sup> A module on management of snake bite cases for Medical Officers, 2015 published

by the Government of West Bengal is in practice in Alipurduar and Medical Officers have been trained on snake bite management based on it with re-orientations every year<sup>6</sup>.

Referral rate is low given the fact that neither CCU nor facility for dialysis was available during the period under reference. The death rate due to snakebite is low compared other parts of the state<sup>14</sup>. Now, with the provision of a Critical Care Unit (CCU) and a Dialysis unit at the Alipurduar District Hospital, Snake bite management has been strengthened.

Our study covers only the Alipurduar District Hospital and not any other government hospital in the district where snakebite management is provided and hence does not reflect the true load of snake bite cases in government hospitals in Alipurduar. Further, as this study was hospital based, it does not encompass the significantly large group of patients who did not report to the district hospital for various reasons. However, it does give a trend of the epidemiological and clinical profile of snake bite cases in the district. A community-based approach would have overcome this limitation and provided a true picture of the magnitude of snake bites in the district along with the detailed epidemiological and clinical profile of the snake bite cases in the district. Further, a prospective hospital based study with a written protocol would have provided better epidemiological and clinical profile of snake bite cases, factors associated with adverse outcomes and provided information on strengthening snake bite management in the district.

#### CONCLUSION

Snake bite, a neglected tropical disease with high fatality, can be treated effectively based on syndromic approach as per the state snake bite management protocol. AVS without skin test dose is life saving and is very safe and effective with Injection Adrenalin at hand. Medical Officers need to be trained and regular re-orientation on usage of the snake bite management protocol done to ensure better detection of signs of envenomation and judicious use of AVS. Timely hospitalization, avoidance of traditional or faith healers and increased awareness among the masses with regards to prevention of snake bites, including use of personal protection during outdoor activities, deterrence to attraction of snakes to households by improving storage of food and fodder at home and avoiding keeping or rearing of fowls at homes or in the vicinity, clearing garbage and vegetations from peri-domestic areas, sleeping on charpoys and using mosquito nets while sleeping. Apart from community awareness on prevention of snake bites paramedical staff and the community at large needs to be sensitized on proper and appropriate first aid in cases of snake bites, including avoidance of ligatures, need for

psychological assurance to the snake bite victim and rapid transportation to the nearest hospital where AVS is available. Lastly, a community based survey to assess the magnitude and epidemiological profile of snake bites in the district, as well as, a prospective hospital based study to study the clinical profile of snake bite cases along with study of factors associated with adverse outcomes with a view of improve clinical management of snake bites in the district is recommended.

**Patients' consent :** No direct consent was taken from the patients as this is a retrospective study. Details of history, clinical findings, admissions and outcomes were obtained from the hospital records. Consent, instead was taken from the Superintendent of the hospital to use information contained in the patient records without disclosing the identity of the patients and solely for educational and research purposes with the view to decreasing snake bite morbidity and mortality.

**Financial support and sponsorship :** Nil

**Conflicts of interest :** There are no conflicts of interest

#### REFERENCES

- Warrel DA, Snake bite — *Lancet* 2010; **375(9708)**: 77-88.
- Jean-Philippe Chippaux. Snakebite envenomation turns again into a neglected tropical disease! *Journal of Venomous Animals and Toxins including Tropical Diseases* 2017; **23**: 38.
- Mohapatra B, Warrell DA, Suraweera W, Bhatia P, Dhingra N, Jotkar RM, *et al* — Snakebite Mortality in India: A Nationally Representative Mortality Survey. *PLoS Negl Trop Dis* 2011; **Apr 12**: **5(4)**: e1018
- Govt of India, Central Bureau of Health Intelligence, Health Status Indicators, National Health Profile. 2007 and 2008 (Provisional): 3.1.2.9 State/UT wise Cases and Deaths due to Snake bite in India. 103-18. <http://cbhidghs.nic.in/writereaddata/mainlinkFile/Health.StatusIndicators.pdf> (accessed 2011)
- Hati AK, Mandal M, De MK, Mukherjee H, Hati RN — Epidemiology of Snake bites in the district of Burdwan, West Bengal. *J Indian Med Assoc* 1992; **90(6)**: 145-7.
- Public Health Branch of the Directorate of Health Services, Government of West Bengal. A Module on Management of Snake bite cases for Medical Officers, 2015
- Raja R, Dasgupta N, Mistry N, Chowdhury A — Glimpses of Ophiofaunal diversity in Jalpaiguri district of West Bengal, India. *International Journal of Physical and Social Sciences* 2013; **3**: Issue 11.
- Simpson ID — A study of the current knowledge base in treating snake bite amongst doctors in the high-risk countries of India and Pakistan: does snake bite treatment training reflect local requirements? *Trans R Soc Trop Med Hyg* 2008; **102(11)**: 1108-14. doi: 10.1016/j.trstmh.2008.04.013. Epub 2008 May 27.
- Ian D — Simpson. A study of current knowledge base in treating snake bite among doctors in the high risk countries of India and Pakistan: does snake bite treatment training reflect local requirements? *Transactions of the Royal Society of Tropical Medicine and Hygiene* 2008; **102**: 1108-14.
- Pakistan Medical Research Council. Endorsed by World Health Organisation. Snake bite management in Asia and Africa, A guide to snake bite in the key areas for mortality & morbidity.
- Sharma SK, Chappuis F, Jha N, Bovier PA, Loutan L, Koirala S — Impact of Snake bites and determinants of fatal outcomes in South Eastern Nepal. *Am J Trop Med Hyg* 2004; **71(2)**: 2234-38.
- Indrani D, Mandal P, Mahata R, Ghosh S, Ghosh B, Das R — Clinical and Epidemiological Profile of Snake bite cases in a Tertiary Care Medical College and Hospital in Eastern India. *Saudi J Med Pharm Sci* 2017; **3(5)**: 408-13.
- Pal A, Mukherjee S — Clinical profile and outcome of Snake bite in patients attending a Tertiary Care Hospital in Bankura, West Bengal. *IOSR Journal of Dental and Medical Sciences*-eISSN: 2279-0853, p-ISSN: 2279-0861. Volume 17, Issue 10 Ver. 7 (October 2018) PP 34-39
- Majumdar D, Sinha A, Bhattacharya SK, Ram R, Dasgupta U, Ram A — Epidemiological Profile of Snake Bite in South 24 Parganas District of West Bengal with focus on Underreporting of Snake bite deaths. *Indian Journal of Public Health* 2014; **58(1)**: 17-21.
- Guidelines for the Management of Snake Bites. 2<sup>nd</sup> Edition. WHO 2016
- Islam AT, Sultana S, Uddin MK, Rahman M — Snakebites with Neuropsychiatric Presentation – A Study in the Hill Tracts of Bangladesh. *Journal of Enam Medical College* 2018; **8(1)**:
- Williams SS, Wijesinghe CA, Jayamanne SF, Buckley NA, Dawson AH, Laloo DG, *et al* — Delayed Psychological Morbidity Associated with Snakebite Envenoming. *PLoS Negl Trop Dis* 2011; **5(8)**: e1255.
- Warrel DA — Guidelines for the management of snake bites. WHO 2010. ISBN 978-92-9022-377-4.
- Chew SK, Khor HW, Ahmad R, Hisamuddin N, Abdul Rahman N — A five-year retrospective review of snakebite patients admitted to a tertiary university hospital in Malaysia. *International Journal of Emergency Medicine* 2011, **4**: 41.
- Sarojini C, Jhansi Rani C, Mukherjee PK, Chandramouli V — Clinico-epidemiological profile and outcome of snakebite victims in South West of West Bengal: A District hospital based study. *Perspective in Medical Research*, May-August 2018; **6(2)**:
- Pandey DP — Epidemiology of Snake bites Based on Hospital Survey in Chitwan and Nawalparsai Districts, Nepal. *Journal of Nepal Health Research Council* 2005; **4(2)**:
- Koushik M, Ghosh R, Gantait K, Saha K, Parua P, Chatterjee U, *et al* — Incidence and treatment of snakebites in West Bengal, India. *Toxicology Reports* 2019; **6**: 239-43.
- Sharma N, Chauhan S, Faruqi S, Bhat P, Varma S — Snake envenomation in a North Indian hospital. *Emerg Med J* 2005; **22(2)**: 118-120. (PMC free article PubMed)



## Original Article

# A comparative study of Internal Jugular Vein Cannulation : Ultrasound Guided Technique *versus* Conventional Blind technique

Vaibhavi B Patel<sup>1</sup>, Smita R Engineer<sup>2</sup>, Kiran B Patel<sup>3</sup>, Vikram P Mehta<sup>4</sup>, Mona Bilandani<sup>5</sup>

Central line cannulation is usually done in right Internal Jugular Vein (IJV) for long term administration of fluid, antibiotic, hemodialysis, chemotherapy. IJV cannulation is routinely done by Conventional Blind Technique (CT) using landmarks. Nowadays Ultrasound Guided Technique (UT) is used to cannulate IJV. The aim of this study was to compare outcome of the Ultrasound guided technique with conventional technique in terms of success rate, number of attempts for IJV location and cannulation, time required for procedure and complications. This study was conducted on 120 patients who were randomly divided into 2 groups. For both the group of patients seldinger technique was used to cannulate IJV. In Ultrasound guided technique, IJV was cannulated with real time linear transducer probe with high frequency. IJV was cannulated in first attempt in 90% of patients with Ultrasound guided technique and 61.66% of patients with conventional technique (P=0.03). Second attempt was required in 10% of patients with Ultrasound guided technique and 30% of patients with conventional technique (P=0.01). Third attempt was required in 8.33% of patients with conventional technique while it was nil in Ultrasound guided technique. Mean time for procedure was 7.78+2.75 minutes in Ultrasound guided technique and 19.4+5.68 minutes in conventional technique (p=0.025 is significant). In Ultrasound guided technique, rate of complications like, carotid puncture was none, hematoma 8%, difficulty in cannulation 1.66% which are less as compared to conventional technique (8%, 21.66%, 10% respectively). Thus, use of real time Ultrasound for IJV cannulation is beneficial than conventional technique in terms of decreased number of attempts for IJV location and cannulation, less access time and decreased incidence of complications.

[J Indian Med Assoc 2020; 118(1): 17-21]

**Key words :** IJV cannulation, Ultrasound Guided Technique, Conventional Technique.

Central line cannulation is usually done in Internal Jugular Vein (IJV) because of its anatomical position, larger diameter and minimal obstruction in its route to right atrium. IJV cannulation is commonly performed to obtain access to central venous line for long term administration of fluid, antibiotic, total parenteral nutrition, hemodynamic monitoring (CVP-central venous pressure monitoring), hemodialysis, chemotherapy<sup>1</sup>. Cannulation of right IJV has been the preferred method for central venous cannulation<sup>2</sup> because right IJV is in a straight line to right atrium and shorter route, while on the left side the dome of the pleura is higher and chances of damage to thoracic duct. IJV cannulation is done routinely by conventional blind technique using landmarks. Nowadays, Ultrasound Guided Technique is used to cannulate IJV. Ultrasound is a non-invasive, non-ionising form of imaging that is safe for all

age group patients. It has been suggested that Usg Guided Technique has higher success rate, less number of attempts for IJV location and cannulation, less time required for IJV cannulation and less procedure related complications<sup>3</sup>. Failure to successfully cannulate IJV occurs in the range of 2% to 9% in conventional technique<sup>4</sup>. The aim of this study was to compare outcome of the Ultrasound Guided Technique for IJV cannulation with Conventional Technique in terms of number of attempts for IJV location and cannulation, total access time (time from skin puncture to aspiration of venous blood) and complications related to the procedure.

### MATERIAL AND METHOD

This study was conducted in Civil Hospital, Ahmedabad on randomly selected 120 patients who required IJV cannulation. It is a prospective, randomized, interventional, controlled type of study. CTRI registration of the study was not done. All the patients were evaluated before the procedure and written informed consent was taken. Patients of age group 18-75 years, with normal coagulation profile and platelets count >80,000 were included in our study. Patients with abnormal coagulation profile, known

Department of Anaesthesia and Surgery, B J Medical College, Civil Hospital, Ahmedabad 380016

<sup>1</sup>MD (Anaesthesia), Tutor and Corresponding Author

<sup>2</sup>MD (Anaesthesia), Associate Professor

<sup>3</sup>MD (Anaesthesia), Assistant Professor

<sup>4</sup>MS (Surgery), Assistant Professor of Surgery

<sup>5</sup>MBBS, First year Resident in Anaesthesia

vascular abnormality were excluded from the study. IJV cannulation was done in various patients undergoing for major prolonged surgery (pheochromocytoma, whipple's surgery, large thyroid mass, neurosurgery), intraoperative CVP monitoring, patients requiring long term antibiotics/fluid, patients requiring TPN (total parenteral nutrition), patients undergoing for hemodialysis, in patient with difficult peripheral venous access. Procedure was carried out in operation theatre in full aseptic precaution with necessary monitors applied and equipments available. No intravenous medication was given before or during procedure. Patients were randomly divided into 2 groups of 60 each, group conventional technique (CT) and Ultrasound Guided Technique (UT) using computer generated random list. In conventional technique group, IJV was cannulated by landmark guided seldinger technique. In Ultrasound Guided Technique group, IJV cannulation was done by using Ultrasound imaging system (SONOSITE) with a 7.5-MHz transducer probe connected to a two-dimensional ultrasound device. The patient was kept in supine position with head turned to opposite side for complete exposure of procedure side and a shoulder pack was placed to extend the neck (except in patient of cervical spine injury in whom the neck was kept in supine position only). After applying monitors (SPO<sub>2</sub> and ECG), painting and drapping was done and Inj. Lignocaine 2% was injected on the site of needle entry.

In conventional technique surface landmarks are apex of the triangle made by 2 heads (sternal and clavicular) of sternocleidomastoid muscle and medial third of clavicle just lateral to carotid pulsations. 20 Gauge needle was advanced at 45 degree angle to skin directed towards ipsilateral nipple. After aspiration of free flow of venous blood in saline filled syringe, a guide wire was passed through the needle into IJV and needle was removed. Finally 18 G radio-opaque catheter was cannulated over guide wire into IJV and guide wire was removed. Catheter was secured with proper stitches.

In Ultrasound Guided Technique, we used real time (Dynamic) method which allows needle observation as the needle approaches the target vessel. The linear transducer probe with high frequency was used, since the vascular structures are typically shallow. Transducer probe was kept in sterile plastic bag with ultrasonic gel applied on it. We kept the probe perpendicular to the floor which is parallel and approximately 1.5 cm above the clavicle (Outplane technique), so only tip of advancing needle can be seen on monitor. Position of IJV in relation to carotid artery was accessed first and 2% lignocaine was injected with 24 g hypodermic needle 0.5 cm above the site of probe. The wheel in subcutaneous tissue was visualized on monitor as enlarging hypoechoic area. 20 G needle was advanced through this skin wheel towards IJV keeping vein in centre

of the monitor. Tip of needle (visualized as hyperechoic dot) was advanced towards anterior wall of IJV. The vein was pierced by a short stabbing motion of needle which leads to puncture in anterior wall of IJV without causing double wall puncture. When the tip of the needle was located in vein, confirmation was done by aspiration of venous blood. We also confirmed IJV location by keeping the probe in plane of IJV so we were able to visualize anterior and posterior walls of vein with tip of needle inside. Then insertion of 18 G catheter was done by seldinger's technique same as mentioned in conventional technique. Successful placement of catheter was confirmed by Chest X-ray and post-procedure arterial blood gas analysis.

In our study, data recorded were patients' demographic data, number of attempts for IJV location, total access time (time from skin puncture to aspiration of venous blood) and complications related to procedure (such as hematoma, arterial puncture, difficulty in IJV cannulation, hemothorax, pneumothorax, brachial plexus irritation, failure of procedure). Primary outcome of this study is successful cannulation in first attempt and secondary outcome is duration of procedure and complications related to procedure.

We have calculated sample size by using power analysis and sample size version 8(2008) software. According to this, sample size was calculated to be 58 in each group at 80% power of study and 0.05 level of significance ( $\alpha$ ) using two-tailed chi-square test. It was done to detect absolute difference of 20% for successful cannulation of IJV in first attempt by Ultrasound Guided Technique and conventional technique. To be on safer side, we decided to keep 60 patients in each group.

Statistical package for social science version 23 (23, IBM, Chicago, USA) was used for analysing the data. All data are presented as mean and % of patients. A chi-square test was used to compare categorical variables and a student's t-test was used to compare independent means. P value <0.05 was considered significant.

#### OBSERVATION

Our study was a single blinded observational type of study carried out on 120 patients who required IJV cannulation. Our 2 study groups were similar in regards to patients' demographic data (age, sex, weight) and site of IJV insertion (Tables 1 & 2).

IVJ was cannulated in first attempt in 90% of patients with Ultrasound Guided Technique and 61.66% of patients with conventional technique ( $p=0.03$ ). Second attempt was required in 10% of patients with Ultrasound Guided Technique and 30% of patients with conventional technique ( $p=0.01$ ). Third attempt was required in 8.33% of patients with conventional technique while it was nil in Ultrasound Guided Technique (Table 3).

Table 1 — Demographic Data		
Group	Usg guided technique (n=60)	Conventional technique (n=60)
Age (Years)	40.31+14.84	43.95+17.97
Sex (M/F)	36/24	23/37
Site of IJV (Right/Left)	59/1	57/3

Table 2 — Number of attempts for successful vein localization			
No of patients cannulated in	UTN (%)	CTN (%)	P Value
First Attempt	54(90%)	38(61.66%)	0.03
Second Attempt	6(10%)	18(30%)	0.01
Third Attempt	0(0)	5(8.33%)	

Table 3 — Total Time Required For IJV Cannulation			
Mean time for IJV Cannulation (minutes)	Usg guided technique	Conventional technique	P value
	7.78+2.75	19.4+5.68	0.025

Total access time is considered from skin puncture to aspiration of venous blood. Mean time for procedure was 7.78+2.75 minutes in Usg Guided Technique and 19.4+5.68 minutes in Conventional Technique having P=0.025 which is significant (Table 4).

In Ultrasound Guided Technique there were no cases of carotid artery puncture while it occurred in 8% of patients in Conventional Technique. Hematoma occurred in only 8% of patients with Ultrasound Guided Technique as compared to 21.6% of patients in Conventional Technique. Difficulty in threading guidewire after successful IJV cannulation occurred only in 1.6% of patients with Usg guided technique as compared to 10% of patients in Conventional Technique. There were no cases of haemothorax, pneumothorax, and brachial plexus injury in any of techniques during the study.

**Ultrasound Guided Images — Outplane technique :** The probe is kept perpendicular to the floor which is parallel and approximately 1.5 cm above the clavicle. In this technique tip of advancing needle and adjacent vascular structure is seen on monitor (Figs 1 & 2).

**Inplane technique :** Probe is kept perpendicular to clavicle and in line with anatomical location of IJV. Whole course of advancing needle towards vein can be seen on monitor (Figs 3 & 4).



Fig 1




Fig 2

Table 4 — Showing Complications		
Complications	Usg guided technique	Conventional technique
Carotid puncture	0(0)	5(8%)
Hematoma	5(8%)	13(21.66%)
Difficulty in cannulation	1(1.66%)	6(10%)




Fig 3




Fig 4

### DISCUSSION

As per our study, in Ultrasound Guided IJV cannulation procedure time (7.78+2.75 minutes in UT) is less compared to Conventional Technique (19.4+5.68 minutes in CT), procedure related complications are negligible with use of Ultrasound, number of attempts for IJV cannulation are less with Ultrasound Guided Technique. (IJV was cannulated in first attempt in 90% of patients with Ultrasound Guided Technique and 61.66% of patients with Conventional Technique)

Considerable amount of time and efforts are spent on insertion of central venous catheter for various purposes, but IJV cannulation is not without risk and may lead to adverse effects that are both hazardous to patients and expensive to treat. Nowadays Ultrasonography is most widely used imaging technique in medical practice. USG is portable, nonradiating, and inexpensive as compared to other imaging modalities such as MRI, CT scan. In modern Ultrasound, a small pulse of echo is transmitted from transducer to body which penetrates body tissue along the path of transmission. Some waves are reflected back to transducer and some continue to penetrate deeper depending upon different acoustic impedances of tissues. The wavelength and frequency of Ultrasound are inversely related. Superficial structures are visualized with high frequency waves and low frequency waves are used to visualize deeper structures<sup>5</sup>. Ultrasonography has specific advantages in cannulating a vessel over conventional blind technique like visualization of a vascular structure, optimal needle placement, avoids puncture of posterior wall of IJV, accurate placement of catheter inside IJV, confirmation of guidewire inside IJV, less number of attempts and lower rate of complications.

As the IJV is a superficial vascular structure, the linear transducer probe with high frequency was used. Most of the patients in our study showed lateral and anterolateral lying IJV relative to carotid artery. We measured depth of IJV from skin which was approximately 0.8+0.4 cm (Fig 5).

On radiological visualization artery is circular in shape, pulsatile, thick walled and non-compressible when pressure is given by probe. While vein is elliptical in shape, non-pulsatile, thin walled and compressible by pressure.

Ultrasonography is useful for central line cannulation as either dynamic technique or static technique. In dynamic technique (also known as real time), the whole procedure is accompanied by the use of USG<sup>6</sup>. While in static technique, USG is used pre-procedure for marking the puncture site, visualization of anatomy and post-procedure to detect possible complications. We used dynamic approach for our study as it allows real-time visualization of the needle tip placement and has been shown to be superior to the static approach in most situations. There are 3 different approaches for Ultrasound Guided Technique for IJV cannulation like outplane, inplane and oblique<sup>7,8</sup>. Outplane (transverse axis) approach has advantage of visualization of adjacent structures and short learning curve but disadvantage of higher risk of injury to the posterior wall of the vessel. Inplane (longitudinal axis) approach has advantage of visualization of vessel, guide wire, catheter during whole procedure but disadvantage is inability to visualize adjacent structures. Third approach is, the oblique, is an intermediate to the previous two<sup>9</sup>. In our study, we kept the probe perpendicular to the floor and parallel to the clavicle (Outplane technique), so only tip of advancing needle and adjacent structures can be seen on monitor. Later on, after aspiration of venous blood, we confirmed IJV location by keeping the probe perpendicular to clavicle (inplane of IJV), so we were able to visualize anterior and posterior walls of vein with tip of needle inside.

All datas are normally distributed in both the Ultrasound Guided Technique and Conventional Technique groups in terms of age, gender (m/f) and site of IJV insersion (rt/lr).

A Study on 200 patients done by Azmat Riaz et al described that vein was localised on 1<sup>st</sup> attempt in 99% of ultrasound-group patients and in 89% of land-mark-group patients. 2<sup>nd</sup> attempt was required in 1% of patients of ultrasound-group and 7% of patients of land-mark-group<sup>10</sup>. In our study, IJV was cannulated in first attempt in 90% of patients with Ultrasound Guided Technique and 61.66% of patients with Conventional Technique (P=0.03). Second attempt was required in 10% of patients with

Ultrasound Guided Technique and 30% of patients with Conventional Technique (P=0.01). Third attempt was required in 8.33% of patients with Conventional Technique while it was nil in Ultrasound Guided Technique




Fig 5

(Table 2). Thus study results are comparable with our study.

Carotid arterial puncture is the most common and frequent complication of IJV cannulation due to its anatomical proximity of 2 vessels. According to our study, in Ultrasound Guided Technique there were no cases of carotid artery puncture while it occurred in 8% of patients with Conventional Technique. Hematoma occurred in only 8% of patients with Ultrasound Guided Technique as compared to 21.6% of patients with Conventional Technique (Table 4). As per study by Darko Sazdov, Marija Srceva *et al* carotid artery puncture occurred in 1% of patients of ultrasound group, while in 8% of patients in land-mark technique (p=0.0007). Hematoma occurred in 4% of patients with Ultrasound as compared to 10% of patients with landmark technique (p=0.01)<sup>10</sup>. This data are comparable with our study results. Similar results were found by Gurkan Turker *et al* which shows carotid puncture and hematoma are less with Usg (0.5% and 1% respectively) as compared to conventional technique (4.73% and 3.68% respectively)<sup>11</sup>. The rate of complications such as accidental puncture of the carotid artery, hemothorax, pneumothorax, and hematoma, as well as the total procedure duration and the occurrence of bloodstream infection associated with catheter, was significantly lower in the group that underwent Ultrasound Guided catheterization<sup>12</sup>. Superiority of Ultrasound Guided Technique over Conventional Technique is seen in patients with obesity, oedema, coagulation disorders, difficult anatomical landmarks, chest deformity, previous surgery, burns, cervical spine injury where neck movements are restricted, hypovolemic patients. Due to proven benefits of Ultrasound Guided IJV cannulation, it will be difficult to justify not using Usg for this procedure. The National Institute for Clinical Excellence (NHS 2002) guidelines issued in 2002 recommend the use of ultrasound for IJV cannulation in both adults and children in most circumstances in both elective and emergency situations<sup>13</sup>. In February 2012, International Committee of Ultrasound

Vascular Access published its recommendations and concluded that, the vascular catheterization must be guided by Ultrasound due to its efficacy and safety provided by the procedure. The National Institute of Clinical Excellence (NICE) issued guidelines recommending that Ultrasound Guidance be used for all elective and emergency central venous cannulations in the National Health Service (England and Wales)<sup>14</sup>.

However there are limiting factors like unavailability of Ultrasound machine due to high cost and need for an experienced operator who has an understanding of probe orientation, image display, the physics of ultrasound, and mechanisms of image generation and artefacts. In our study, Out of total 120 patients, in 2 patients of burns contracture involving neck and in 2 patients with cervical spine injury in whom neck movements were restricted, Ultrasound Guided Technique was used because conventional technique guided IJV cannulation was not possible at all. There were 3 cases in which IJV cannulation was failed by blind technique and later on succeeded by Ultrasound Guided Technique. So success rate of IJV cannulation was 100% with Ultrasound Guided Technique and 95% with conventional technique. Similar results were observed in study done by Shah H, Bhavsar M<sup>15</sup>. Thus use of USG for IJV cannulation reduces mechanical and infectious complications<sup>16</sup>, reduces chances of thrombosis, increases success rate and decreases patient discomfort.

#### CONCLUSION

Use of real time Ultrasound for Internal Jugular Vein cannulation is beneficial than conventional landmark guided technique in terms of decreased number of attempts for IJV location and cannulation, decreased access time, decreased incidence of complications, increased success rate and substantially decreased patient discomfort. Patient discomfort was not measured using any scoring system, but patients in whom IJV cannulation was done with Ultrasound were more comfortable in terms of short procedure time, less number of attempts and less complications.

#### REFERENCES

- 1 Maecken T, Grau T — Ultrasound imaging in vascular access. *Crit Care Med* 2007; **35(5 Suppl)**: S178-85.
- 2 Pat O Daily, Randall B Griep, Shumway NE — Percutaneous internal jugular vein cannulation. *Arch Surg* 1970; **101**: 1103-8.
- 3 Leung J, Duffy M, Finckh A — Real-time ultrasonographically guided internal jugular vein catheterization in the emergency department increases success rates and reduces complications: a randomized, prospective study. *Ann Emerg Med* 2006; **48(5)**: 540-7.
- 4 McGee DC, Gould MK — Preventing complications of central venous catheterization. *N Engl J Med* 2003; **348(12)**: 1123-33.
- 5 Lawrence JP. Physics and instrumentation of ultrasound. *Crit Care Med*. 2007; **35(8 Suppl)**: S314-S322.
- 6 Dowling M, Jjala H, Hardman JG, Bedforth NM — Real-time three-dimensional ultrasound-guided central venous catheter placement. *Anesth Analg* 2011; **112(2)**: 378-81.
- 7 Stone MB, Nagdev A, Murphy MC, Sisson CA — Ultrasound detection of guidewire position during central venous catheterization. *Am J Emerg Med* 2010; **28(1)**: 82-4.
- 8 Phelan M, Hagerly D — The oblique view: an alternative approach for ultrasound-guided central line placement. *J Emerg Med* 2009; **37(4)**: 403-8.
- 9 Ultrasound Guided Internal Jugular Venous Cannulation: Comparison with Land-Mark Technique Azmat Riaz, Rao Ali Shan Khan and Faisal Salim Journal of the College of Physicians and Surgeons Pakistan 2015, **25(5)**: 315-9.
- 10 Darko S, Srceva MJ, Todorova ZN — Comparative analysis of Ultrasound guided central venous catheterization compared to blind catheterization: Contributions. Sec. of Med. Sci., 10.1515/prlozi-2017-0028
- 11 Turker G, Kaya FN, Gurbet A, Aksu H, Erdogan C, Atlas A — Internal jugular vein cannulation: an ultrasound-guided technique versus a land-mark-guided technique. *Clinics (Sao Paulo)* 2009; **64(10)**: 989-92.
- 12 Karakitsos D, Labropoulos N, De Groot E, Patrianakos AP, Kouraklis G, Poularas J — Real-time ultrasound-guided catheterisation of the internal jugular vein: a prospective comparison with the landmark technique in critical care patients. *Crit Care* 2006; **10(6)**: R162.
- 13 Daniel Hind, Neill Calvert, Richard McWilliams, Andrew Davidson, Suzy Paisley, Catherine Beverley, et al. Ultrasonic locating devices for central venous cannulation - Meta analysis. *BMJ* 2003; **327**: 361-4.
- 14 Calvert N, Hind D, McWilliams RG, Thomas SM, Beverley C, Davidson A — The effectiveness and cost-effectiveness of ultrasound locating devices for central venous access: a systematic review and economic evaluation. *Health Technol Assess* 2003; **7(12)**: 1-84.
- 15 Shah H, Bhavsar M, Ultrasound-guided internal jugular venous cannulation comparison with palpation technique: benefits and drawbacks; *IJRRMS* 2013; 3(4)
- 16 O'Grady NP, Alexander M, Burns LA, Dellinger EP, Garland J, Heard SO, et al — Healthcare Infection Control Practices Advisory Committee (2011) Guidelines for the prevention of intravascular catheter-related infections. *Am J Infect Control* 2011; **39(Suppl 1)**: S1-34.

## Original Article

## Prevalence of Autoantibodies in patient complaining of multiple joint pain in a tertiary care hospital

Sangeeta Das Ghosh<sup>1</sup>, Mohiruddin Sk<sup>2</sup>, Sandip Ghosh<sup>3</sup>, Indrani Bhattacharyya<sup>4</sup>

Autoimmune phenomenon is attributed to a number of diseases which were once considered idiopathic. In humans, production of auto antibodies (a-Abs) against self-antigens is quite frequent but earlier their presence was associated with autoimmune diseases, however a-Abs have been documented in non-autoimmune disorders i.e. complicated pregnancy, cancer, stroke etc. This study was designed to determine serum level of antinuclear antibody (ANA), Rheumatoid Factor (RF) and anti dsDNA antibodies in apparently healthy population with multiple joint pain. After written informed consent, blood sample of 294 subjects was obtained by random sampling. Participants of established autoimmune diseases were excluded. Enzyme linked immune sorbent assay (ELISA) was used to determine ANA, RF and anti-dsDNA antibody. Categorical variables were compared by using  $\chi^2$  test. A p value <0.05 was considered statistically significant. Rheumatoid factor was the most frequent a-Ab (19.05%), followed by anti dsDNA (7.14%), while ANA was the lowest (3.4%) antibody detected. Only RF had a statistically significant association with gender ( $p=0.047$ ). No association of these antibodies with age was detected. Rheumatoid factor auto antibody was more prevalent as compared to ANA and dsDNA antibody in healthy adults.

[J Indian Med Assoc 2020; 118(1): 22-4]

**Key words :** ANAs, Anti dsDNA, Anti RF, Autoantibodies, Autoimmune disease.

Anti-nuclear antibodies (ANA) are immunoglobulin directed against autologous cell nuclear and cytoplasmic components<sup>1-3</sup>. The occurrence of different ANA is associated with autoimmune disease and with differences in disease severity including extent of skin involvement, internal organ manifestation and prognosis<sup>2</sup>. Researchers have been performing steady efforts to develop tests for detecting ANA and disease-specific auto antibodies to nuclear antigens for the diagnosis, prognostic assessment, and monitoring of patients with systemic autoimmune diseases<sup>4</sup>. Nowadays, measurement of ANA has been widely used to provide supporting evidence of a diagnosis of autoimmune disease such as Systemic Lupus Erythematosus (SLE), Sjögren etc<sup>5</sup>. SLE is a multisystem disorder that is considered as a prototype Immune Complex (IC)-mediated disease<sup>6</sup>. This autoimmune disease related to central or peripheral nervous system; about 17% to 75% of patients respectively<sup>7</sup>. Additionally, levels of antibodies

against dsDNA were shown covary with SLE disease activity<sup>8</sup>. The aim of this study was to investigate the prevalence of ANA and anti-dsDNA in patient with suspected autoimmune disease.

Rheumatoid Factor (RF) is heterogeneous antibody of IgM class; directed against Fc fragment of IgG. It is used as a disease marker of Rheumatoid Arthritis (RA)<sup>9</sup> but it can be detected in other connective tissue and inflammatory disorders. About 1-5% of healthy individuals may have this antibody and they are at increased risk to develop RA<sup>10</sup>.

## MATERIALS AND METHODS

Serum samples were obtained from different Department of Calcutta National Medical College and Hospital. Each of these serum samples was tested for the presence of ANA, RA factor and anti-dsDNA (Aeskulisa dsDNA check, Aesku Diagnostics, Germany) by ELISA method. These tests were performed by commercial kits according to the manufacturer's instructions. First of all, results were classified as ANA positive or negative according to the definitions contained within the packages for each kit. Subsequently, anti-dsDNA results were classified as positive or negative for each patient. Borderline results were arbitrarily classified as positive.

For statistical analyses, manufacturer suggested cut-off were applied to create positive and negative values from the continuous original observations. Positivity rates,

Department of Microbiology, Calcutta National Medical College, Kolkata 700014

<sup>1</sup>MBBS, MD (Microbiology), Demonstrator

<sup>2</sup>MBBS, MD (Microbiology), Senior Resident and Corresponding Author

<sup>3</sup>MBBS, MS (Orthopaedics), Professor, Department of Orthopaedics, MSVP Calcutta National Medical College, Kolkata 700014

<sup>4</sup>MBBS, MD (Microbiology), Associate Professor

specificities and Spearman correlation coefficient between assays were calculated as indicated using SAS software, Version 9.2 of the SAS system for Windows. In statistical analyzes, p-value <0.05 was considered as significant.

**RESULTS**

In this study, there were more female (214) as compared to male (80). Mean ±SD of age of males and females was 42.20 (3.52) and 24.54 (0.76) ranging from 13-66 and 3-65 (years) respectively. ANA was detected in 4.67% females and 0% males and RF was detected in 10% males and 22.43% females while anti-dsDNA was detected in 5% males and 7.94% females and on comparison there was no significant difference in these parameters. Frequency of ANA, RF and anti-dsDNA was 3.4%,19.04% and 7.14% respectively (Tables 1 & 2).

More females (86%) compared to males (14 %) had a-Abs. Mean age of males was 35.28 years and of females it was 22.14 years. Among the subject 36.74% were more than 35 years (55.1% males and 44.9% females), 50% were between 26-35 years (70.4% males and 29.6% females) and only 13.26% were less than 25 years of age (30.7% males and 69.3% females).

Among the subjects 10 (3.4%) had ANA, 56 (19.05%) had RF and 21 (7.14%) had dsDNA. On comparison of gender, out of 80 males, 8 % had RF, 5% had dsDNA and none of the male had ANA whereas out of 214 females, 22.42% had RF, 7.94% had dsDNA and 4.67% female had ANA. RF was significantly associated with gender (Odds ratio 1.948, 95% confidence interval, (1.004-3.785) while ANA and dsDNA were not associated with gender. It was observed that more females than males (22.42% versus 8%) had RF. Further, none of the a-Abs was associated with age (Figs 1 & 2).

**DISCUSSION**

In the present study, ANA was detected in 3.4% of healthy individuals with multiple joint pain which is lower




Fig 1 — Male ANA, RF and Anti-dsDNA test result Comparison Chart




Fig 2 — Female ANA, RF and Anti-ds DNA test result Comparison Chart

Variable	Male (n=80)	Female (n=214)	Total (n=294)	p- values
ANA Positive	0	10	10	0.068
RF Positive	8	48	56	0.044
Ani-dsDNA Positive	4	17	21	0.6097

Variable	Age group		p value
	<25 (n=39) Positivity (n)%	>25 (n=255) Positivity (n)%	
ANA Positive	1	9	1
RF Positive	12	44	0.1456
Anti-dsDNA Positive	5	16	0.1906

than already documented ie, 4% to 13%<sup>11-14</sup>. In the current study, there was high prevalence of RF in females (22.43%) than males (10%)<sup>15</sup>. It should be noted that RF a-Abs are frequently present in healthy subjects and can be detected in chronic infections. ANA positivity rate found in our female patients is a consistent result with the knowledge of the autoimmune diseases are more frequent in women<sup>16,17</sup>. This predominancy was researched by Leo and et al. According to their study, the hormone profile, fetal microchimerism and some strategic genes which are on the sex chromosomes are playing role on this relationship<sup>18</sup>. Our three years' experience of testing autoantibodies was shared in this study. Reliable test results are very important for the health of the patients with autoimmune disorders. For being a dependable laboratory, having enough knowledge and experience about the chosen methods of autoantibody tests is mandatory. A good relationship with the clinicians is also an indispensable component of confidential analysis and reporting.

### CONCLUSION

Rheumatoid factor auto antibodies were more prevalent as compared to dsDNA and ANA in healthy adults. Further, RF was associated with gender as it was prevalent more in males compared to females. ANA and anti dsDNA were not associated with age and gender.

### Conflict of interest :

None of the researchers has any financial or other interest in the products that were used for this study.

### ACKNOWLEDGMENTS

We acknowledge the financial support of MSVP of CNMCH and technical and logistic support of HOD Microbiology. We are also grateful to all the subjects who donated blood for this study.

### REFERENCES

- Cepeda EJ, Reveille JD — Autoantibodies in systemic sclerosis and fibrosing syndromes: clinical indications and relevance. *Curr Opin Rheumatol* 2004; **16(6)**: 723-32.
- Mierau R, Moinzadeh P, Riemekasten G, Melchers I, Meurer M, Reichenberger F — Frequency of disease-associated and other nuclear autoantibodies in patients of the German network for systemic scleroderma: correlation with characteristic clinical features. *Arthritis Res Ther* 2011; **13(5)**: R172.
- Walker JG, Fritzler MJ — Update on autoantibodies in systemic sclerosis. *Curr Opin Rheumatol* 2007; **19(6)**: 580-91.
- Kavanaugh A, Tomar R, Reveille J, Solomon DH, Homburger HA — Guidelines for clinical use of the antinuclear antibody test and tests for specific autoantibodies to nuclear antigens. American College of Pathologists. *Arch Pathol Lab Med* 2000; **124(1)**: 71-81.
- Tan EM, Cohen AS, Fries JF, Masi AT, McShane DJ, Rothfield NF — The 1982 revised criteria for the classification of systemic lupus erythematosus. *Arthritis Rheum* 1982; **25(11)**: 1271-7.
- Koffler D, Agnello V, Thoburn R, Kunkel HG — Systemic lupus erythematosus: Prototype of immunocomplex nephritis in man. *J Exp Med* 1971; **134(3)**: 169-79.
- Bonfa E, Golombek SJ, Kaufman LD, Skelly S, Weissbach H, Brot N, *et al* — Association between lupus psychosis and anti-ribosomal P protein antibodies. *N Engl J Med* 1987; **317(5)**: 265-271.
- Mok CC, Lau CS. Pathogenesis of systemic lupus erythematosus. *J Clin Pathol* 2003; **56(7)**: 481-90.
- Hugle B, Hinze C, Lainka E, Fischer N, Haas JP — Development of positive antinuclear antibodies and rheumatoid factor in systemic juvenile idiopathic arthritis points toward an autoimmune phenotype later in the disease course. *Pediatr Rheumatol Online J* 2014; **12**: 28.
- Alam SM, Kidwai AA, Jafri SR, Qureshi BM, Sami A, Qureshi HH, *et al* — Epidemiology of rheumatoid arthritis in a tertiary care unit in Karachi, Pakistan. *J Pak Med Assoc* 2011; **61(2)**: 123-6.
- Baig MM, Shere SJ — Prevalence of autoantibodies in Saudi population. *J Med* 1989; **20(3-4)**: 286-90.
- Manoussakis MN, Tzioufas AG, Silis MP, Pange PJE, Goudevenos J, Moutsopoulos HM — High prevalence of anti-cardiolipin and other autoantibodies in healthy elderly population. *Clin Exper Immunol* 1987; **69(3)**: 557-65.
- Goemaere S, Ackerman C, Ghoethals K, De Keyser F, Vander Straeten C, Vebruggen G, *et al* — Onset of symptoms of rheumatoid arthritis and relation to age, sex and menopausal transition. *J Rheumatol* 1990; **17(12)**: 1620-2.
- Azizah MR, Shahnaz M, Zulkifli MN, Nasuruddin BA — Antinuclear, anti-mitochondrial, anti-smooth muscle and anti-parietal cell antibodies in the healthy Malaysian population. *Malaysia J Pathol* 1995; **17(2)**: 83-6.
- Mordvinov GV, Mordvinova IV. Prevalence of rheumatoid factor in the healthy population of Moldova Republic. *Klin Lab Diagn* 2000; **(12)**: 33-5.
- Güdücüoğlu H, Yaman G, Çikman A, Çalısır U, Berktaş M — Retrospective evaluation of immunoblotting (IB) test results in anti-nuclear antibody positive patients. *Turkish Journal of Clinical Laboratory* 2011; **2**: 59-62.
- Li Q, Karp D, Quan J, Branch VK, Zhou J, Lian Y, *et al* — Risk factors for ANA positivity in healthy persons. *Arthritis Res Ther* 2011; **13(2)**: R38.
- Lleo A, Battezzati PM, Semli C, Gershi ME, Podda M — Is autoimmunity a matter of sex? *Autoimmun Rev* 2008; **7(8)**: 626-30.

*If you want to send your queries and receive the response on any subject from JIMA, please use the E-mail or Mobile facility.*

### **Know Your JIMA**

Website	: <a href="http://www.ejima.in">http://www.ejima.in</a> <a href="https://onlinejima.com">https://onlinejima.com</a>
For Editorial	: <a href="mailto:jima1930@rediffmail.com">jima1930@rediffmail.com</a> Mobile : +919477493027
For Circulation	: <a href="mailto:jimacir@gmail.com">jimacir@gmail.com</a> Mobile : +919477493037
For Marketing	: <a href="mailto:jimamkt@gmail.com">jimamkt@gmail.com</a> Mobile : +919477493036
For Accounts	: <a href="mailto:journalacct@gmail.com">journalacct@gmail.com</a> Mobile : +919432211112



## Original Article

## Cross sectional analysis of suicidal behavior in adolescents — comparison with adult attempters

P N Suresh Kumar<sup>1</sup>, P K Anish<sup>2</sup>

Indian studies have suggested that the suicides and suicidal behaviours are higher in younger age groups. Studies comparing the socio-demographical and clinical variables between various age groups in relation to suicidal behaviours are required to tailor specific interventions for the early age groups. Hence this study was conducted to analyse socio-demographic and clinical variables of adolescent suicide attempters and to compare certain potential risk factors between adolescent and adult suicide attempters. Socio-demographic and clinical variables of 199 adolescents versus 960 adult attempters were compared using a specially designed questionnaire to find out the potential risk factors in adolescents. Significant number of adolescent attempters were females, from nuclear family, had higher frequency of medical illnesses and specific life stressors related to their developmental period. Majority had medical contact within one week of attempt and attempted suicide within one week after the stressor. Current psychiatric diagnosis was more frequent in adults than adolescents. In adolescents the most frequent psychiatric diagnosis was adjustment disorder. This study concludes that Adjustment disorders related to life stressors preceded adolescent suicidal attempts more frequently than adults. Helping adolescents to cope with life events should be a focus of suicide prevention strategies in this population.

[J Indian Med Assoc 2020; 118(1): 25-9]

**Key words :** Adolescents, suicide, attempts, life events, self-harm.

Worldwide, suicide is the second leading cause of death in young people, representing a major public health problem especially in the 15-19 year age group<sup>1</sup>. Youth is a period of heightened risk of suicide and suicide is a leading cause of death among young people in India<sup>2</sup>. A cross sectional study of adolescent suicidal behaviours (age 12-19) in two schools in Delhi found the life time prevalence of suicidal attempts to be as high as 8%<sup>3</sup>. Adolescents suicide attempters are a heterogeneous group and their characteristics are different and the pattern of suicidal behaviour is not similar to that of adult population<sup>4</sup>. Potential risk factors for suicide attempts in adolescents include female gender, psychopathology especially a major depressive disorder, previous suicide attempts, hopelessness, recent stressful life events, suicide attempts by family members or friends, chronic physical illness, family violence and dysfunction and lower academic achievement<sup>5</sup>. Suicidal attempt is the strongest risk factor for suicide, psychiatric disorders which tend to onset in adolescence being the second<sup>6</sup>. Other contributors include genetic vulnerability, psychological, familial, social, and

cultural factors<sup>7</sup>.

The precipitating events, which have led to a suicide attempt, are most often stressful interpersonal problems between the adolescent and his parents or peers<sup>8</sup>. Some types of stress such as exit events, or interpersonal losses, and other major negative events often precede suicide attempts<sup>9</sup>. The relationship between chronic stress and adolescent suicide attempts is clinically important because treatment for disharmony among parents may reduce the risk of suicide among adolescents living in a stressful home environment<sup>10</sup>. Given the potentially tragic nature of adolescent suicide attempts and the elevated risk of suicide clustering among adolescents, the identification of adolescents at risk for suicide attempts before their behaviour escalates and becomes more serious would be of obvious value.

There is a lack of systematic research on suicide in adolescence. Due to paucity of data and to increase our understanding of adolescent suicide attempt and improve the management present study was conducted with the following aims:

- (1) To study the socio-demographic and clinical variables of adolescent suicide attempters.
- (2) To compare certain potential risk factors between adolescent and adult suicide attempters (Tables 1-3).

### MATERIAL AND METHODS

A non-experimental descriptive approach was used for

Department of Psychiatry, KMCT Medical College, Calicut, Kerala 673602

<sup>1</sup>MD, DPM, DNB, PhD, MRC Psychiatry, Professor and Corresponding Author

<sup>2</sup>MD, Assistant Professor, Department of Psychiatry, Institute of Mental Health and Neurosciences, Calicut 673008

Table 1 — Socio-Demographic Details

	Age below 19 years (199)	Age above 19 years (960)	p
<b>Sex :</b>			
Male	76 (38.2)	501 (52.2)	0.00*
Female	123 (61.8)	459 (47.8)	
<b>Marital Status :</b>			
Unmarried	174 (87.4)	272 (28.3)	0.00
Married	24 (12.1)	641 (66.8)	
Separated/widow	1 (0.5)	47 (4.9)	
<b>Religion :</b>			
Hindu	136 (68.4)	762 (79.4)	0.00
Muslim	59 (29.6)	156 (16.3)	
Christian	4 (2.0)	40 (4.2)	
Tribal	0 (0.0)	2 (0.2)	
<b>Domicile :</b>			
Rural	161 (80.9)	833 (86.8)	0.02*
Urban	38 (19.1)	127 (13.2)	
<b>Mean Education (Years)</b>	9.41±2.31	7.65±3.65	0.00
<b>Occupation-Unemployed</b>	155 (77.9)	441 (45.9)	0.00
<b>Family :</b>			
Joint	62 (31.2)	413 (43.1)	0.007
Nuclear	137 (68.2)	547 (56.9)	
<b>Consanguinity- yes</b>	16 (8.0)	63 (6.6)	0.45*
<b>Medical Illnesses</b>	39 (19.6)	29 (3.0)	0.001*
<b>F/H/O Pseh Illness</b>	59 (29.6)	332 (34.6)	0.19*
<b>F/H/O suicide</b>	5 (2.5)	69 (7.2)	0.01*
<b>Past Psych Illness</b>	22 (11.1)	249 (25.9)	0.00*
<b>Past Suicide attempt</b>	23 (11.6)	194 (20.2)	0.002*
<b>Mean Number of past attempts</b>	0.13±0.5	0.24±0.7	0.004

\*Fisher's exact test

Table 2 — Special Causes for the Attempt

	Age Below 19 years (199)	Age Above 19 years (960)	p
Special cause for attempt	171 (85.9)	827 (86.1)	0.51
Friction with family members	74 (37.2)	130 (13.6)	0.00
Love failure	19 (9.5)	40 (4.2)	0.00
Death of dear ones	14 (7.0)	28 (2.9)	0.01
Friction with girl/boy friends	11 (5.5)	15 (1.6)	0.00
Failure in Exam	11 (5.5)	0 (0.0)	0.00
Physical problems	7 (3.5)	66 (6.9)	0.05

\*Fisher's exact test

this study. The study was conducted at KMCT Medical College Hospital with an inpatient capacity of 60 beds. Psychiatry Department receives referrals from various Departments for management of approximately 10-20 suicide attempters in any given month. The institution has a written rule that any patient admitted for a suicide attempt has to be evaluated by the Psychiatry Department before his or her discharge from the hospital. Consecutive suicide attempters of all age groups referred for detailed psychiatric evaluation from various Departments from the period 2012 January to 2015 December formed the study sample for the two comparison arms. Information was obtained from patient, care giver and other reliable sources. The assessments were completed within one day of receiving

Table 3 — Details about Suicide Attempt

	Age below 19 years (199)	Age above 19 years (960)	p
<b>H/O Medical Contact :</b>			
Within 1 week	162 (81.4)	685 (71.2)	0.04
Within 1 week to 1 month	13 (6.5)	78 (8.1)	
Within 1 month to 3 months	13 (6.5)	105 (10.9)	
<b>Hours of Sleep :</b>			
0-2	5 (2.5)	52 (5.4)	0.00
3-5	46 (23.1)	502 (52.3)	
6 or more	142 (71.3)	378 (39.3)	
<b>Change in Weight :</b>			
Nil	185 (93.0)	759 (79.0)	0.00
10% or Weight Loss	14 (7.0)	200 (20.8)	
10% or more Weight gain	0 (0.0)	1 (0.0)	
<b>Time Difference between stress and Attempt :</b>			
24 Hours	89 (44.7)	188 (19.6)	0.00
24 Hours – 1 week	27 (13.6)	111 (11.6)	
1 week – 1 month	20 (10.1)	98 (10.2)	
More than 1 month	40 (20.2)	483 (50.3)	
<b>Suicide Treats</b>	84(42.2)	438(45.6)	0.21*
<b>Suicide Notes</b>	21 (10.6)	85 (8.9)	0.42*
<b>Time of Suicide Attempt :</b>			
12 Midnight- 6am	8 (4.0)	46 (4.8)	0.97
6am – 6 pm	108 (54.3)	519 (54.1)	
6pm – 12 Midnight	80 (40.2)	382 (39.8)	
<b>Place of Attempt :</b>			
House	156 (78.4)	745 (77.6)	0.95
Outside	40 (20.1)	202 (21.0)	
<b>Consumption of Alcohol at the Time of Attempt</b>	11 (5.5)	211 (22.0)	0.00
<b>Mode of Attempt :</b>			
Poisoning	147 (73.9)	757 (78.9)	0.21
Drowning	0 (0.0)	9 (0.9)	
Hanging	24 (12.1)	97 (10.1)	
Cutting	28 (14.1)	97 (10.1)	
<b>Mean Time of Discovery (Minutes)</b>	77.4+140.6	73.5+152.8	0.74
<b>Mean time of Reaching Hospital (Minutes)</b>	198.0+485.0	167.0+304.9	0.39

\*Fisher's exact test

referral. World Health Organisation (WHO) definition of child and adolescent such as any person between ages 10 and 19 was adopted for the selection of adolescents<sup>11</sup>. The total sample was divided into two groups, Group 1, 10-19 years age and group 2, above 19 years age (Table 4).

#### INCLUSION AND EXCLUSION CRITERIA

Suicide attempters who are willing to participate in this study after signing the written informed consent were included in the study. The subjects had to be inpatients during the assessment and should be able to respond to the tools appropriately. Subjects who had no accompanying family member for corroboration of information were excluded.

#### TOOLS

A self-designed socio-demographic questionnaire was used to document the psycho-socio-demographic details and life events. It consists of a total of 104 items covering various aspects like socio-demographic data, details of

Table 4 — *Psychiatric Diagnosis*

	Age below 19 years (199)	Age above 19 years (960)	p
Psychiatric Diagnosis			
- Present	120 (60.3)	764 (79.6)	0.00
Depression	22 (11.1)	261 (27.2)	0.00
Mania/Bipolar	2 (1.2)	29 (3.0)	0.78
Schizophrenia/Psychoses	2 (1.2)	53 (5.5)	0.00
Alcohol/Drug abuse	7 (3.5)	117 (12.2)	0.00
Adjustment Disorder	81 (40.7)	324 (33.8)	0.04
Neuroses	21 (10.6)	65 (6.8)	0.05
Fisher's exact test			

the suicide attempt, past medical and psychiatric history, current psychiatric diagnosis and psychosocial events. Initially a pilot assessment was conducted on 10 subjects and some modifications were done in the questionnaire like removing the ambiguity of some items. Test retest reliability of the prepared tool was established by Cronbach's alpha (0.714). Psychiatric diagnosis was based on DSM IV Criteria<sup>12</sup>.

Detailed explanations were given to subjects and parents about the purpose of the study. Confidentiality of the information was assured and informed consent was taken from parents prior to enrolling subjects. Rapport was established and explanation was given about the study tool. Total time taken for data collection was one hour for each subject.

#### ANALYSIS OF DATA

Data was analyzed by SPSS-10 PC software system. Socio-demographic data and Psychological factors were analyzed by percentage of frequencies. Association between socio-demographic and selected psychological factors between two groups was analyzed using "t" test, "Chi-square" test and "Fisher's exact test".

#### RESULTS

Table 1 shows the comparison of socio-demographic variables of adolescent versus adults suicide attempters. Significant number of adolescent attempters were females, from nuclear family and had history of medical illnesses. Adult attempters had higher frequency of family history of suicide, past psychiatric illnesses, past suicide attempts and mean number of attempts.

Table 2 shows the comparison of causes for the attempt. Adolescents had significantly higher frequency of events such as friction with family members, love failure, death of loved ones, friction with girl/boyfriends, failure in exam, physical problems etc.

Table 3 shows that significantly more adolescents had medical contact within 1 week of attempt and attempted suicide within 1 week after the stressor. Table 4 shows that current psychiatric diagnosis was more frequent in adults than adolescents. The most frequent psychiatric diagnosis in adolescents was adjustment disorder.

#### DISCUSSION

Suicidal behavior in adolescent age group is an important public health issue because of the large proportion (21.8%) that 10-19 year olds constitute in the population of India<sup>13</sup>. In a study which evaluated the cause of death among those aged 10-19 years, in a rural population of 108,000 in South India, suicide accounted for about a quarter of all deaths in males and between 50% and 75% of all deaths in females aged 10-19 years. The average suicide rate for girls was 148 per 100,000, and for boys, 58 per 100,000<sup>14</sup>. In the adolescent age group, the associated developmental and social challenges are quite different from that of an adolescent population. Understanding the characteristics and variables associated with adolescent suicide in contrast to adult suicidal behaviours is important in designing interventions specifically tailored to this population.

In most of the earlier studies, girls outnumbered boys in suicide attempt and ideation<sup>15,16</sup>. Present study also there was an over representation of female attempters though it was not clinically significant. An earlier study on adolescent suicides from India has shown equal representation of male and female attempters<sup>17</sup>. This gender paradox is explained by recent studies<sup>18</sup>. Adolescent females feel shy in discussing their views and problems with peers or families resulting in feeling of being isolated and attempt suicide as a cry for help. Furthermore, although adolescent boys have less suicidal thoughts, their ideas translate into more completed suicides. Other possibility for this difference could be due to a cohort effect (ie, more recent female generations are engaging in more suicidal behavior). Additionally, boys may not remember or may minimize the clinical significance of this type of behavior during their adolescence (ie, retrospective reporting grossly underestimates the rates of adolescent suicidal behavior).

Though both the groups had predominance from rural areas (reflecting the catchment area of the hospital), there was a significant difference between the groups with a larger proportion of adolescents from urban areas. Majority of adolescents were found to be hailing from nuclear families. This could be related to fact that most of these adolescents are from urban settings where joint family fragmentation is more likely. Further studies are required to explore the psychiatric morbidity and suicide ideation among the adolescents living in different social settings like rural joint families and urban nuclear families. It also needs to be explored whether there is an underreporting of adolescent suicides in rural settings due to cultural factors.

Majority of adolescent subjects attempted suicide within 24 hours after the stressor whereas the adult subjects attempted more than a week after the stressor. This may suggest that the adolescent attempts were more

unplanned and impulsive. This again ties up with the fact that adolescent suicides appear to be related more with situational problems and associated concerns about social support and coping abilities.

Studies have found that suicidal people frequently consult medical services, usually a primary care physician, shortly before the act, sometime in the last few days or hours prior to the suicidal act<sup>19</sup>. In the present study 94% of the adolescents had contact with medical personnel at least three months before the attempt. Among them, 81% had medical contact within one week of their attempt. Even though the information is not exactly available from the subjects whether they had suicidal ideation during these consultations, considering the predominance of physical and Psychological problems in them, it is a strong possibility having grave implications. One investigation found that 41% of adult persons who committed suicide had contact with a health care professional within 28 days of death, 47 percent within 1 week, and 18 percent on the day of death. However, suicidal intent was discussed in only 22% of this cases<sup>20</sup>. Unfortunately similar data is not available from adolescent group. Educational measures for primary care physicians aimed at identifying the flag signs of suicidality will be useful adolescent suicide prevention.

Many physicians miss or dismiss the telltale signs of suicidality presuming that it is quite unlikely to occur in adolescents. They also have the feeling that asking suicidal ideation may provoke suicidality in certain situations. This lack of recognition and misbelief is compounded by the fact that suicidal intent often presents differently in adolescents and can be confused with changes in personality, sleep problems, drug abuse, fatigue, concentration deficits, diminished memory, lack of initiative etc. They are also less likely to report suicidal ideation and intent compared to adults. This makes difficulty in establishing the diagnosis and initiating treatment. The problem is further compounded by the lack of sensitization among primary care givers regarding adolescent mental health problems. Hawton et al reported that only 20% of their adolescent study sample had Psychiatric disorders<sup>21</sup>. In the present study 60% had psychiatric disorders but with a predominant diagnosis of adjustment disorder. Their attempts were mostly impulsive occurring within a week after the stressor. This again underlines the fact that adolescent suicides appear to be related more with poor social support and maladaptive coping techniques.

In the adult suicidology, one avenue to understand the psychological condition immediately prior to the self-destructive act has been the study of suicide notes. In a study by Posener et al in Montreal, seventeen adolescents who left notes were identified, comprising 10% of the population of suicides<sup>22</sup>. Victims who left notes did not differ from the total group in age and sex distribution. In

the present study also only 10% had written suicide notes prior to their attempt. Even though this is a small number, suicidal messages from children and adolescents need to be taken very seriously as it sometimes may offer an opportunity for suicide prevention.

In contrast to adult counter parts, the adolescent group had lesser rates of past psychiatric illnesses, past suicidal attempts, family history of psychiatric illnesses/suicide and current psychiatric illnesses. These findings shows that adult attempters may have more severe psychiatric problems compared to adolescents where attempts are more likely to be related to immediate life stressors. These findings have important implications in developing specific suicide prevention strategies in adolescents. Adequate focus on screening for adjustment problems and developing healthy coping strategies as well as ensuring availability of social support for handling psycho-social stressors should be given due importance in adolescents.

Psychoactive substance abuse is a major problem in adolescents especially so in those who attempt suicide. However, in the present study only a small proportion had a history of alcohol or drug abuse and intoxication at the time of attempt. Other than adjustment disorder, neurotic disorders also significantly outnumbered among adolescents.

Stressors such as conflicts with parents, breakup of a relationship, school difficulties or failure, death of dear ones, and physical ailments were the commonly cited reasons by adolescents for attempting suicide in this study. Among young people, suicidal behavior was found to be associated with female gender, not attending school or college, independent decision making, premarital sex, physical abuse at home, lifetime experience of sexual abuse, and probable common mental disorders. Violence and psychological distress were independently associated with suicidal behavior. Factors associated with gender disadvantage increased vulnerability, particularly in rural women<sup>23</sup>. Family disruptions and discord stemming from excessive arguments and overt violence, loss of relatives due to marital separation or divorce, and problems in family interpersonal relations are reported as stressful circumstances experienced by suicidal adolescents<sup>24</sup>. We also found out that the quality of stressful life events the adolescents have faced in the preceding month is significantly different from adults. This particular aspect highlights the need of early intervention focusing on specific issues in the adolescent period related to their developmental period.

Categorizing the stress according to the source may also be useful. Majority of the adolescents in our study had interpersonal problems with family members and friends as the main precipitating factor for the attempt. When adolescents have problems in their close relationships with

family and friends, they may lose important sources of social support which may in turn increase the risk of suicidal behaviour. Early intervention may be necessary to protect the quality and integrity of these interpersonal relationships. The occurrence of these stressful life events in the recent past should be "red flags" for the clinician working with potentially suicidal adolescents. However, it is also important to note that these events most often occur without suicide as a consequence.

#### LIMITATIONS

Our study does have certain limitations. Cross sectional nature of the study, possibility of retrospective bias especially in the aftermath of the suicide attempt, lack of scales for assessing psychiatric morbidity or severity of suicidal risk could be the important ones. However, considering that the study is a non-purposive comparison study, any rater bias appears to be unlikely. Another issue would be the comparatively smaller sample size in the adolescent group. But the strength of our study is that all the evaluations have been consistently done by the same person (a trained psychiatrist) on a consecutive sample.

#### CONCLUSIONS

This study clearly differentiates adolescent suicide attempters from adult counter parts with certain unique risk factors. Adolescent attempters were mostly females, hailing from nuclear urban family, had higher frequency of medical illnesses and had specific life events related to their developmental period. Majority had medical contact within one week of attempt and attempted suicide within a week after the stressor. Their most frequent psychiatric diagnosis was adjustment disorder. Adult attempters were mostly males, had higher frequency of family history of suicide, past psychiatric illnesses, past suicide attempts and current psychiatric diagnoses.

This study suggests psychosocial intervention with specific focus on developmental issues as the most important strategy for adolescent suicide prevention. Future research should examine the role of those intervening variables that are capable of reducing the negative impacts of chronic strains, such as social supports, personality styles and coping skills. Studies with control groups comprising non-suicidal adolescents are also necessary for confirmation of our findings. Adolescents experiencing higher rates of cumulative stressful life events should be the target population for repeated monitoring for identification of suicidal behaviour.

#### REFERENCES

- 1 Hawton K, Saunders KE, O'Connor RC — Self-harm and suicide in adolescents. *Lancet* 2012; **379(9834)**: 2373-82.
- 2 Vijayakumar L, John S, Pirkis J, Whiteford H. Suicide in developing countries (2): Risk factors. *Crisis* 2005; **26(3)**:112-9.
- 3 Sidharta T, Jena S — Suicidal behaviors in adolescents. *Ind J Pediatrics* 2006; **73(9)**: 783-88.

- 4 McClure GMG — Suicide in children and adolescents in England and Wales 1960-1990. *Br J Psychiatry* 1994; **165(4)**: 510-14.
- 5 Lewinsohn PM, Rhode P, Seeley JR — Psychosocial risk factors for future adolescent suicide attempts. *J Consulting & Clinical Psychology* 1994; **62(2)**: 297-305.
- 6 Kessler RC, Berglund P, Demler O, Jin R, Merikangas KR, Walters EE — Lifetime prevalence and age-of-onset distributions of DSM-IV disorders in the National Comorbidity Survey Replication. *Arch Gen Psychiatry* 2005; **62(6)**: 593-602.
- 7 Nock MK, Borges G, Bromet EJ, Cha CB, Kessler RC, Lee S — Suicide and suicidal behavior. *Epidemiol Rev* 2008; **30**: 133-54.
- 8 Cohen-Sandler R, Berman AL, King RA — Life stress and symptomatology, determinants of suicidal behaviour in children. *J Am Acad Child Adol Psychiatry* 1982; **21(2)**: 178-86.
- 9 Slater J, Depue RA — The contribution of environmental events and social support to serious suicide attempts in primary depressive disorder. *J Abnorm Psychol* 1981; **90(4)**: 275-85.
- 10 Adams DA, Overholser JC, Spirito A — Stressful life events associated with adolescent suicide attempts. *The Canadian Journal of Psychiatry / La Revue canadienne de psychiatrie* 1994; **39(1)**: 43-48.
- 11 <https://www.britannica.com/science/adolescence>
- 12 American Psychiatric Association - Diagnostic and Statistical Manual of Mental Disorders. 4<sup>th</sup> ed. Text Revision (DSM-IV-TR). Washington, DC: American Psychiatric Association; 1994.
- 13 Government of India. Census of India 2011 - Accessed on 20<sup>th</sup> August 2018. [censusindia.gov.in/2011-Documents/Houselisting%20English.pdf](http://censusindia.gov.in/2011-Documents/Houselisting%20English.pdf)
- 14 Aaron R, Joseph A, Abraham S, Mullyil J, George K, Prasad J, et al — Suicides in young people in rural southern India. *Lancet* 2004; **363(9415)**: 1117-8.
- 15 Garfinkel BD, Froese A, Hood J — Suicide attempts in children and adolescents. *Am J Psychiatry* 1982; **139(10)**: 1257-61.
- 16 Lewinsohn PM, Rhode P, John R — Adolescent Suicidal Ideation and Attempts: Prevalence, Risk Factors, and Clinical Implications. *Clin Psychol Sci Prac* 1996; **3**: 25-46.
- 17 Kumar CT, Chandrasekaran R — A study of psychosocial and clinical factors associated with adolescent suicide attempts. *Ind J Psychiatry* 2000; **42(3)**: 237-42.
- 18 Thakur D, Gupta A, Thakur A, Mazta SR, Sharma D — Prevalence and predictors of suicidal ideations among school going adolescents in a hilly state of India. *Ind J Psychiatry* 2015; **24(2)**: 140-3.
- 19 Emanuel EI, Fairclough DL, Slutsman J, Emanuel LL — Understanding economic and other burdens of terminal illness: the experience of patients and caregivers. *Ann Intern Med* 2000; **132(6)**: 451-56.
- 20 Goodwin R, Olfson M — Self-perception of poor health and suicidal ideation in medical patients. *Psychol Med* 2002; **32(7)**: 1293-9.
- 21 Hawton K, O'Grady J, Osborn M, Cole D — Adolescents who take overdoses, their characteristics, problems and contacts with helping agencies. *Br J Psychiatry* 1982; **140**: 118-23.
- 22 Posener JA, LaHaye A, Cheifetz PN — Suicide notes in adolescence. *Can J Psychiatry* 1989; **34(3)**: 171-6.
- 23 Pillai A, Andrews T, Patel V — Violence, psychological distress and the risk of suicidal behaviour in young people in India. *Int J Epidemiol* 2009; **38(2)**: 459-69.
- 24 Pfeffer CR — Suicidal behavior in children and adolescents: causes and management. In: Lewis M Editor - Child and adolescent Psychiatry — a comprehensive textbook. 3rd Edition. Philadelphia: Lippincott Williams & Wilkins, 2002; 796-805.

## Original Article

## Prevalence of anaemia in school going adolescents in a municipality town of West Bengal

Ritesh Singh<sup>1</sup>, Banasree Bhadra<sup>2</sup>, Soumyajit Moulik<sup>3</sup>, Suman Kumar Roy<sup>4</sup>

Anaemia is a major public health problem in India. It directly causes many premature deaths in population. The deaths due to anaemia as indirect cause far outweigh the number due to many well-known causes of mortality in a developing nation. A cross-sectional survey was conducted in two schools of a municipal town of West Bengal. One school was public and the other was private. The haemoglobin was estimated by HaemoCue® instrument. Two hundred students finally participated in the study. 118 (59%) of these students were studying in a government run institute. 105 (53%) of the study participants were females. The mean (SD) age of the study participants was 14.05 (0.8) years. Majority of the participants belonged to lower middle class. About half (51%) of the study participants have correct knowledge about the cause of anaemia. Out of three major meals of the day, study participants were taking dinner regularly. Around 47% of them missed or never took a breakfast within last week. Around 91% of the study participants were non-vegetarian, a usual diet pattern in this part of the country. The mean (SD) Haemoglobin concentration of students was 12.58 (1.20) gm/dl. The mean haemoglobin concentration of male students was significantly more than that of female. Overall 70 (35%) students were suffering from anaemia. 45 out 105 (42.9%) female students were suffering from anaemia, whereas 25 (26.3%) male students were suffering from anaemia. Male students were significantly more tall and heavier when compared with female students. Anaemia was statistically more common in students belonging to low socioeconomic strata of the society.

The awareness about anaemia and its causes is low in school students. The prevalence of anaemia is high amongst school going adolescents in urban areas of West Bengal.

[J Indian Med Assoc 2020; 118(1): 30-4]

**Key words :** Anaemia, Prevalence, India, Adolescents, Diet, Nutrition.

Anaemia is the qualitative and or quantitative diminution of haemoglobin or red blood cells (RBC) or both in respect to the age and sex of individual. The net effect of which is failure in transport of adequate amount of oxygen to the tissues. Generally, anaemia is said to occur when Haemoglobin (Hb) is less than a critical value which depends upon the sex and age of the individual, but symptoms depend not only on the reduction but also on the rate of reduction of Hb<sup>1</sup>. There are several different types of anaemia and each one has a different cause. Nutritional anaemia of which iron deficiency anaemia is a

sub-type is the most common type. Other nutritional causes of anaemia are lack of vitamin B12 or folic acid. Iron deficiency anaemia occurs when there is lack of iron in the body. The major dietary sources of iron are meat, dried fruit and some vegetables. Some food items and medicines make iron absorption difficult. Iron is used by the body to make haemoglobin, which helps store and carry oxygen in red blood cells. There are many conditions that can lead to a lack of iron. In men, and postmenopausal women the most common cause is bleeding in the stomach and intestines caused by regular and indiscriminate use of non-steroidal anti-inflammatory drugs (NSAIDs), peptic ulcer, stomach cancer or bowel cancer. In women of reproductive age, the most common causes of iron deficiency anaemia are heavy menstruation and pregnancy. If iron deficiency anaemia is left untreated it can make one susceptible to repeated episodes of illness and infection, as lack of iron in the body affects one's immune system. Severe iron deficiency anaemia may increase the risk of developing complications that affect the heart or lungs such as, tachycardia and heart failure. Pregnant women also have a

Department of Community and Family Medicine, All India Institute of Medical Sciences, Kalyani 741245

<sup>1</sup>MBBS, MD (Community Medicine), Associate Professor

<sup>2</sup>MBBS, MD (Obstet & Gynaecol), FICMCH, FIAOG, Associate Professor, At present : Professor, Department of Obstetrics and Gynaecology, College of Medicine and JNM Hospital, WBUHS, Kalyani 741235, and Corresponding Author

<sup>3</sup>MBBS, Intern, College of Medicine and JNM Hospital, WBUHS, Kalyani 741235

<sup>4</sup>MBBS, MD (Community Medicine), Professor and Head, Department of Community Medicine, College of Medicine and JNM Hospital, WBUHS, Kalyani 741235

higher risk of complications before and after birth. A major underlying cause of maternal deaths in India is anaemia.

Anaemia is India's major public health problem. There is a wide range of prevalence data available from different regions of the country. In India, anaemia is primarily due to poor nutrition. According to the fourth round of National Family Health Survey (NFHS - 4) conducted during 2015-16, anaemia is prevalent in 53% of women in the age group of 15 to 49 years. It is present in 23% of men in the age group of 15 to 45 years<sup>2</sup>.

Adolescence is the period of rapid growth when iron requirement for both boys and girls increase. The incidence of iron deficiency anaemia among adolescents is rising. The awareness regarding anaemia and appropriate diet is extremely poor among adolescents which are made worse by the misleading advertisements in the media. Risk factors for anaemia among adolescents are low intake of meat, fish, poultry and iron fortified foods, frequent dieting, vegetarian life style, heavy menstrual periods, rapid growth, overweight and obesity.

Keeping in view of the importance of anaemia amongst adolescents in Indian context and paucity of studies relating the dietary habits and prevalence of anaemia amongst school going students, a cross-sectional study was done with the aim of finding the prevalence of anaemia in the adolescent boys and girls.

#### MATERIALS AND METHODS

A cross-sectional survey was conducted with the primary objective of finding out prevalence of anaemia amongst school going students. The secondary objectives were finding out the factors associated with decreased level of haemoglobin in school students and finding out the gender difference in haemoglobin level and factors associated with these differences. Both boys and girls students studying in two identified schools in the study area were told about the study. They were enrolled once they gave written informed assent. Students studying in standard IX of these schools were enrolled for the study. For comparison purpose schools catering to two different socio-economic background students were covered. Students suffering from serious medical condition where haemoglobin concentration is known to be low were excluded from the study. A sample size of 200 students was calculated based upon the assumption of prevalence of anaemia to be 50% amongst the study population and an absolute error of 7%. We enrolled 100 students studying in standard IX of each of the selected two schools. There was more than one section of class IX in the two schools. Students were randomly selected from the three sections of class IX of each school. The sampling frame contained

only those students of class IX of all sections present on the day of the interview and blood draw for haemoglobin estimation. The study was conducted in Kalyani which is a municipality town in Nadia District of state of West Bengal. The two schools chosen were Kalyani Experimental Higher Secondary School, a renowned private educational institution, located in the heart of the city and the Kalyani Shikshayatan, run by the Government of West Bengal. The later school normally caters to students belonging to lower socio-economic strata of the society.

The clearance from the Institutional Ethics Committee (IEC) was taken prior to initiation of the study. Students were told in detail about the study and the procedure to be performed for measurement of the haemoglobin level. Since the students were below 18 years of age, written informed consent of their parent was taken. The students were free to opt out of the study at any time they felt like. The confidentiality of the study participants was maintained, and no personal identifying data were collected. Proper dietary advices for correction of anaemia were imparted to all students after the blood collection. The students identified to be anaemic were given health education regarding the prevention and management of anaemia. Those found anaemic were referred to their nearest health care centre for further evaluation and management of anaemia.

A semi-structured self-administered questionnaire was designed in English. It had questions related to basic socio-demographics-economic status of the participants. Usual dietary history of students was collected. The questionnaire was pre-tested in the same school but in different class which was not part of the study before the conduction of the study. The haemoglobin of the students was measured by an instrument called HaemoCue<sup>®</sup>. The HemoCue<sup>®</sup> haemoglobin analyser is a portable, rapid and accurate method of measuring haemoglobin at the bedside. It has a sensitivity of 99.4% and specificity of 88.4%<sup>3</sup>. Height and weight of the students were measured following standard procedures. For the study purpose, anaemia was defined as per the World Health Organization criteria of haemoglobin of less than 12 gm/dl in females and males less than 15 years of age. For males of age  $\geq 15$  years the cut-off for anaemia was haemoglobin level of 13 gm/dl<sup>4</sup>.

The data collected were entered in MS Excel 2010. The data was analysed by IBM SPSS<sup>®</sup> version 22.0. The difference of mean haemoglobin concentrations between the schools and also between the genders was tested by independent t- test. Chi-square test was used to compare the proportions. Two-tailed significance test with p value of 0.05 or less was considered to be statistically significant.

## RESULTS

Two hundred students finally participated in the study. 118 (59%) of these students were studying in Kalyani Shikshayatan School, a Government run Institute. 105 (53%) of the study participants were females. The mean (SD) age of the study participants was 14.05 (.8) years. The minimum age was 12 years and the maximum age was 16 years. While 15% of the participants' father possessed some professional qualification, majorly (27.5%) were high school pass. All the study participants were classified for their socioeconomic status using modified Kuppuswamy socioeconomic scale. The income ranges for the year 2019 was taken into account<sup>5</sup>. Majority of the subjects belonged to lower middle class (Table 1).

About half (51%) of the study participants have correct knowledge about the cause of anaemia. Rest were either ignorant or had false information about the causes of anaemia. When asked whether they were suffering from anaemia, 8 (4%) replied affirmatively (Table 2).

Out of three major meals of the day, study participants took dinner regularly. Around 47% of them missed or never took breakfast in the preceding week (Fig 1).

Majority of the students were taking any one or more

Table 1 — Distribution of study participants according to their socioeconomic status (n=200)

Socioeconomic class	Frequency	Percentage
Upper (I)	23	11.5
Upper middle (II)	52	26.0
Lower middle (III)	73	36.5
Upper lower (IV)	52	26.0
Lower (V)	0	0.0
Total	200	100.0

Table 2 — Self-reported prevalence of anaemia (n=200)

Suffering from anaemia	Frequency	Percentage
Yes	8	4.0
No	42	21.0
Don't know	150	75.0
Total	200	100.0




Fig 1 — Intake of regular meals of the day of participants (n=200)

fast foods during 1 to 3 days in a week. 95% of the study participants were in the habit of washing hands before taking any meal. Helminthiasis is one of the prime reasons of anaemia in our country. Students' were asked whether they took an anti-helminthic tablet or suspension within last six months of the date of study. 83% of them either did not take it or they were unaware about it. Around 91% of the study participants were non-vegetarian, a usual diet pattern in this part of the country. Majority (112, 56%) of the non-vegetarians were taking any meat product for 1-3 days in a week. Only 10% of the students were taking any fruit every day. Around 80% of the students were taking at least two servings of green leafy vegetables, a rich source of iron every day. The mean (SD) Haemoglobin concentration of students was 12.58 (1.20) gm/dl. The minimum value of Hb was 9.3 gm/dl and the maximum were 16.4 gm/dl. The mean haemoglobin concentration of male students was significantly more than that of female (13.13 versus 12.09, p value <0.000, Table 3).

Overall 70 (35%) students were suffering from anaemia. 45 out of 105 (42.9%) female students were suffering from anaemia, whereas 25 out of 95 (26.3%) of male students were suffering from anaemia (Table 4). Anaemia was significantly more common in females compared to males (Chi square statistic 5.99, p value: 0.014)

Out of 70, 54 (77%) students were suffering from mild anaemia and 16 (23%) were suffering from moderate anaemia; no one had severe anaemia (Fig 2).

Male students were significantly more tall and heavier when compared with female students (p value <0.01, Table 5).

Anaemia was not statistically associated with the usual food habits of the respondents (chi square statistic 0.454, p value 0.501). It may be due to large number of non-vegetarians in the sample. Anaemia was statistically more common in students belonging to low socioeconomic strata of the society (Table 6).

## DISCUSSION

The findings of this school-based study to find out

Table 3 — Sex-wise distribution of mean haemoglobin of study participant (n=200)

Sex	Mean Hb (in gm/dl)	Standard deviation	t-statistic, p value
Male	13.13	1.07	6.8, <0.000
Female	12.09	1.09	

Table 4 — Prevalence of anaemia in different sexes (n=70)

Sex	Frequency	Percentage	Chi square statistic, p value
Male	25	26.3	5.99, 0.014
Female	45	42.9	
Total	70	35.0	



**Table 5 — Anthropometric detail of study subjects (n=200)**

Sex	Male (n=95)	Female (n=105)	t-statistic and p value
Mean (SD) height in cms	158.8 (9.5)	149.6 (5.3)	8.52, 0.0000
Mean (SD) weight in kgs	46.2 (9.9)	43.4 (7.9)	2.22, 0.027

**Table 6 — Distribution of study participants according to their socioeconomic and anaemia status (n=200)**

Socioeconomic class	No anaemia	Anaemia	Total	Chi square statistic, p value
Upper (I)	16	17	23	10.065, 0.017
Upper middle (II)	42	10	52	
Lower middle (III)	45	28	73	
Upper lower (IV)	27	25	52	
Total	130	70	200	




Fig 2 — Distribution of anaemic students according to the severity of anaemia (n=70)

prevalence of anaemia amongst students revealed that 35% of the total sample population had anaemia. We could not find any case of severe anaemia. Majority were suffering from mild anaemia. Various studies found out the prevalence of anaemia to be between 19 to 88% in different part of the country<sup>6</sup>. Soma Gupta *et al*<sup>7</sup> conducted a study in the Midnapore district of West Bengal wherein they measured the haemoglobin concentration of school students (6 - 16 years). They found out that overall prevalence of anaemia in school going students was 80.2%. It was more in female students (86.1%) compared to male (76.0%). According to a study done by Saratha A *et al*<sup>8</sup> in South India, out of total 300 medical and nursing female students 228 (76%) were anaemic. Of them 170 (56.67%) had mild and 58 (19.33%) had moderate anaemia. A community-based cross-sectional study was carried out in rural areas of West Bengal by Arlappa N *et al*<sup>9</sup> during 2002 - 2003. A total of 437 pre-school children were covered for the estimation of blood haemoglobin levels. Majority (81%) of them were anaemic, and the prevalence was significantly (p<0.001) higher among 1 - 3 year-old (91%) as compared to 4 - 5

year-old (74.6%) children. Santanu K Sharma *et al*<sup>10</sup> did a community-based survey in Assam to find out anaemia amongst adolescent females. They found out that 71.5% of the surveyed females had anaemia. Sabita Basu *et al*<sup>11</sup> conducted a study to assess the prevalence of anaemia among 1120 apparently healthy adolescents (12 to 18 years) sampled from 11 city and 2 rural schools in Chandigarh. All the boys and the girls were subjected to anthropometric examination and haemoglobin estimation. The estimation of haemoglobin was done by cyanmethemoglobin method. The overall prevalence of anaemia calculated as per WHO Guidelines was significantly higher among girls (23.9%) as compared to boys (odds ratio 3.75, 95% CI 2.59 to 5.43, P<0.01). A cross sectional study done in urban schools of Ludhiana, Punjab found out that overall prevalence of anaemia was 51.5%. Girls had a significantly higher prevalence of anaemia except at 5 years and 10-12 years age<sup>12</sup>. Another cross-sectional survey done in Bangalore of South India amongst 2030 boys and girls, aged 5-15 years, attending schools found out that mean (SD) Hb concentration of all children were 12.6 (1.1) g/dl (range 5.6-16.7). The overall anaemia prevalence was 13.6%. Anaemia prevalence was lower in boys than girls (12.0% versus 15.3%)<sup>13</sup>. Researchers estimated the prevalence of iron deficiency anaemia among adolescent schoolgirls in the age group of 13-17 years in Chennai. Haemoglobin was estimated using cyan method. The prevalence of anaemia was found to be 78.75% among school students<sup>14</sup>.

Saratha A *et al*<sup>8</sup> found out that 157 (89.71%) students who did not consume green leafy vegetable regularly were anaemic. They also found out that anaemia was associated with increasing age, increasing academic year, consumption of non-green leafy vegetable and passage of worms in stool. They could not find any significant association between anaemia and consumption of veg/non-veg foods, history of chronic illness, type and duration of menstrual cycle. Similarly, there was no significant association between anaemia with height, weight and BMI. According to Arlappa N *et al*<sup>9</sup> children belonging to lower socioeconomic Scheduled Caste and Scheduled Tribe communities were at higher risk for anaemia (OR=2.3; 95% CI 1.3-3.9). Ludhiana study<sup>12</sup> found out that more menarcheal girls were anaemic as compared to non-menarcheal ones. The prevalence of anaemia was high (38%) even in higher socioeconomic groups. Nearly half (47.6%) of well-nourished children were anaemic. The mean Hb also was lower than expected normal values in both nutritional groups. Compared to non-vegetarians (38%), more vegetarians (65.9%) were anaemic. Chennai study<sup>14</sup> showed significant association of anaemia with type of

family, socioeconomic status and diet. In this study 42.5% of girls with BMI < 18 were found to be anaemic.

The result of the present study is somewhat in between of these extremes. This study is unique in a sense that it was carried out amongst students studying in a particular class of two schools. There was little variability in the age of the sample students. The major difference between the prevalence of anaemia in adolescents in this research and others may be due to three reasons. First, none of the study found out by us was conducted by HemoCue® device. The other reason may be the students were comparatively belonging to better socio-economic strata of the society compared to others studies. The third reason of finding a low prevalence of anaemia may be the school health programme run by the Indian government wherein iron folic acid tablets and albendazole tablet is regularly given to school students. The finding of this study that anaemia is more prevalent in female students compared to male is similar to almost all searched studies. The mean haemoglobin concentration of all students of our study (12.58 gm/dl) is similar to that of the Bangalore study<sup>13</sup> (12.6 gm/dl). We could not find any association between the consumption of vegetarian food with anaemia. Saratha A *et al*<sup>8</sup> also could not find any similar association. Our finding of increased anaemia in lower socioeconomic strata of society is consistent with Arlappa N *et al*<sup>8</sup> study. Low quality of nutrient intake is common in those who cannot afford to buy green leafy vegetable and meat products. The limitation of the study is that the researchers did not attempt to find out the reason of anaemia whether iron, vitamin B12 or folate deficiency in diagnosed cases of anaemia.

We conclude that the awareness about anaemia and its causes is low in school students. The prevalence of anaemia is high amongst school going adolescents in urban areas of West Bengal. Anaemia is more common in female adolescents compare to their male counterparts. Low socioeconomic status is significantly associated with high prevalence of anaemia. The mean haemoglobin concentration of females is low compared to males even when analysed for all children below 15 years of age.

## REFERENCES

- 1 Mahapatra ABS. 2014 — Essentials of Physiology by Mahapatra, Current Books and Publishers.
- 2 International Institute for Population Sciences (IIPS) and ICF. 2017. National Family Health Survey (NFHS-4), 2015-16: India. Mumbai: IIPS.
- 3 Shahshahani HJ, Meraat, N, Mansouri F — Evaluation of the validity of a rapid method for measuring high and low haemoglobin levels in whole blood donors. *Blood transfusion = Trasfusione del sangue*, 11(3): 385–390. doi:10.2450/2012.0031-12.
- 4 World Health Organization (2008). Worldwide prevalence of anaemia 1993–2005. Geneva: World Health Organization. ISBN 978-92-4-159665-7.
- 5 Wani RT. Socioeconomic status scales modified Kuppaswamy and Udai Pareekh's scale updated for 2019. *J Family Med Prim Care* 2019; **8(6)**: 1846-9.
- 6 Srihari G, Eilander A, Muthayya S, Kurpad AV, Seshadri S — Nutritional status of affluent Indian school children: what and how much do we know? *Indian Pediatr* 2007; **44(3)**: 204-13.
- 7 Gupta S, Taraphdar P, Roy TG, Haldar D, Dey SK, Purkait B — The silent burden of anemia in school age children: A community based study in West Bengal. *Indian J Med Sci* 2012; **66(6-8)**: 163-8.
- 8 Saratha A, Singh Z, Datta SS, Boratne AV, Senthilvel V, Joice S — Prevalence of Anaemia among Young Adult Female Students in a Medical Teaching Institution in Pondicherry. *Indian Journal of Maternal and Child Health* 2010; **12(4)**: 1-8.
- 9 Arlappa N, Balakrishna N, Laxmaiah A, Brahmam GNV. Prevalence of anaemia among rural pre-school children of West Bengal, India. *Annals of Human Biology* 2010; **37(2)**: 231-42.
- 10 Sharma SK, Narain K, Devi KR, Mohapatra PK, Phukan RK, Mahanta J — Haemoglobinopathies – major associating determinants in prevalence of anaemia amongst adolescent girl students of Assam, India. *WHO South-East Asia Journal of Public Health* 2012; **1(3)**: 299-308.
- 11 Basu S, Basu S, Hazarika R and Parmar V. Prevalence of Anemia Among School Going Adolescents of Chandigarh. *Indian Pediatrics* 2005; **42(6)**: 593-7.
- 12 Verma M, Chhatwal J, Kaur G — Prevalence of anemia among urban school children of Punjab. *Indian Pediatr* 1998; **35(12)**: 1181-6.
- 13 Muthayya S, Thankachan P, Zimmermann MB, Andersson M, Eilander A, Misquith D, *et al* — Low anemia prevalence in school-aged children in Bangalore, South India: possible effect of school health initiatives. *Eur J Clin Nutr* 2007; **61(7)**: 865-9.
- 14 Premalatha T, Valarmathi S, Sriyayanth P, Sundar JS, Kalpana S — Prevalence of Anemia and its Associated Factors among Adolescent School Girls in Chennai, Tamil Nadu, India. *Epidemiol* 2012; **2**: 2.






Fig 2 — (a) Relation of pseudocyst with stomach, (b) Aspiration to confirm pseudocyst, (c) Cystogastrostomy being fashioned

cells of the pancreas. Trauma is the commonest cause of pseudocyst formation in children (over 60%), when the pancreas gets compressed against the vertebral column. It may also form as a sequelae of pancreatitis, pancreaticobiliary system anomalies like pancreatic divisum, viral illnesses and steroid use. Our patient did not have any history of trauma, nor did her history, biochemical or imaging findings suggest any other aetiology. Hence, it was considered to be idiopathic. The challenge of diagnosing such a case increases, in view of no suggestive aetiological history.

USG is the initial imaging of choice in children, since there is no ionizing radiation. On USG, pseudocysts are seen as unilocular or multilocular (with internal septations), well defined, smooth walled, hypochoic lesions. If hemorrhage or infection is present, internal echoes or may be seen.

CT scan is mandatory for planning the treatment. It identifies the size, margin and nature of the pancreas with presence or absence of peripancreatic fluid. The pseudocyst appears as an area with well defined wall or capsule which enhances on contrast, and a central area of low attenuation. CT scan can accurately evaluate the presence and extent of pancreatic necrosis and peripancreatic fat inflammation.

ERCP / MRCP is essential to rule out any pancreatico-biliary anomalies.

Since paediatric pseudocysts are rare, there are few reported large series, and there is no consensus on optimal management. Pseudocysts <5cm usually resolve spontaneously or with conservative management, which involves hydration, analgesia, bowel rest and parenteral nutrition. Larger cysts are less likely to resolve spontaneously, and may be complicated by rupture, haemorrhage, infection, or lead to gastric outlet obstruction. So, larger pseudocysts and those failing a conservative trial require either internal or external drainage. Asymptomatic cysts >5cm even with minimal symptoms or those without any morphological or size change after a 6 week conservative trial require intervention<sup>1</sup>. Even if drainage is contemplated, conservative management for 4-6 weeks, is needed for cyst wall maturation<sup>2</sup>. Though laparoscopic internal drainage has made advances in management of pseudocyst of pancreas over the last two decades, experience in paediatric population is limited and long-term outcome of relevant studies is awaited<sup>3</sup>. Open internal

drainage, either cystogastrostomy, cystoduodenostomy or cystojejunostomy, remains the mainstay of management, the option depending on the anatomic location of the pseudocyst.

Advances in imaging and endoscopy have made image guided percutaneous drainage and endoscopic drainage safe and effective in skilled hands. Endoscopic treatment aims to create a communication between the cyst cavity and the gastrointestinal tract. However, it is dependent on the anatomical location of the pseudocyst. For endoscopic treatment to be feasible, the distance between the pseudocyst and the gastrointestinal wall has to be <1cm<sup>4</sup>. However, recurrence rates with percutaneous and endoscopic drainage procedures are higher than surgical internal drainage. Since most studies on percutaneous and endoscopic drainage procedures have been performed on adults, further studies to compare these techniques in children are required to determine the optimal management in children<sup>5,6</sup>.

Till then, surgical drainage remains the management of choice in paediatric pseudocyst of pancreas.

**Source of support : None**

**Conflicting Interest : None**

#### REFERENCES

- Bradley EL 3rd — A clinically based classification system for acute pancreatitis. Summary of the International Symposium on Acute Pancreatitis, Atlanta, Ga, September 11 through 13, 1992. *Arch Surg* 1993; **128(5)**: 586-90.
- Khalifa M, Gobran T, Shreef KS Waly A — Pancreatic pseudocyst in children: a single institute experience. *Ann of Pediatr Surg* 2015; **11(2)**: 127-31.
- Zerem E, Hauser G, Loga-Zec S, Kunosia S, Jovanovica P, Crnkic D — Minimally invasive treatment of pancreatic pseudocysts. *World J Gastroenterol* 2015; **21(22)**: 6850-60.
- Rosso E, Alexakis N, Ghaneh P, Lombard M, Smart HL, Evans J, *et al* — Pancreatic pseudocyst in chronic pancreatitis: endoscopic and surgical treatment. *Dig Surg* 2003; **20(5)**: 397-406.
- Makin E, Harrison PM, Patel S, Davenport M - Pancreatic Pseudocysts in Children: Treatment by Endoscopic Cyst Gastrostomy. *J Pediatr Gastroenterol Nutr* 2012; **55(5)**: 556-8.
- Jazrawi SF, Barth BA, Sreenarasimhaiah J — Efficacy of endoscopic ultrasound-guided drainage of pancreatic pseudocysts in a pediatric population. *Dig Dis Sci* 2011; **56(3)**: 902-8.

## JIMA Guidelines for Authors

Communications intended for publication should be sent to the Editor, Journal of the Indian Medical Association (JIMA). JIMA will consider manuscripts prepared in accordance with the **Vancouver style**<sup>1</sup>.

Articles are considered for publication on condition that these are contributed solely to JIMA, that they have not been published previously in print and are not under consideration by another publication. In the selection of papers and in regard to priority of publication, the opinion of the Editor will be final. The Editor shall have the right to edit, condense, alter, rearrange or rewrite approved articles, before publication without reference to the authors concerned.

**Authorship:** All persons designated as authors should **qualify for authorship**. Authorship credit should be based only on **significant contributions** to (a) conception and design, or analysis and interpretation of data; and to (b) drafting the article or revising it critically for important intellectual content; and on (c) final approval of the version to be published. **Conditions (a), (b) and (c) must all be met.** Authors may include explanation of each author's contribution separately.

**Manuscript and CD:** Scripts of articles should be **type-written, double-spaced on one side of A4 size paper with at least 2.5 cm margin at the top and on left hand side of the sheet** and should be submitted in **triplicate**. Also **sending CD in MS Word** is a must. Manuscripts not accepted for publication will not be returned to the authors. Authors should keep one copy of their manuscripts for reference. Separate pages should be used for different sections.

**Title page—**The title page should include the title of the article which should be concise but informative, name(s) of author(s) with his/her (their) academic qualification(s) and designation(s), and complete postal address including pin code of the institution(s) to which the work should be attributed.

**Abstract—**The 2nd page should carry an abstract of no more than 250 words and should contain the purposes of the study or investigations, basic procedure, main findings and their implications along with Key words and Take home message (4-5 bullets).

**Text—**The text of Original Articles should conform to the conventional division of Abstract, Introduction, Material and Method, Observations, Discussion and References. Other types of articles such as Practitioners' Series, Case Reports, Current Topics, GP Forum, etc. are likely to need other formats.

**Statistical evaluation—**Description of the statistical methods used should either be given in detail in the "Material and Method" section of the article or supportive reference may be cited.

**Abbreviations—**Standard abbreviations should be used and **be spelt out when first used in the text**. Abbreviations should not be used in the title or abstract.

**Units of measurement—**Metric units should be used in scientific contributions. If the conventional units or SI units were actually followed in measurements that should be given in parentheses.

**Drugs—**The **generic names of the drugs (and not proprietary names)** including dose(s), route(s) and period of administration should be mentioned.

**Length of manuscripts—**For Originals Articles: Maximum 2000-2200 words, 2-3 figures, and/or 2-4 tables, for Case Reports: Maximum 800 words, 2 figures, 1 table, for Letter to the Editor: 300-500 words.

**Tables—**Tables should be simple, self-explanatory and should

supplement and not duplicate the information given in the text.

**Illustrations—**Graphs, charts, diagrams or pen drawings must be drawn by professional hands in Indian ink (black) on white drawing paper. In case of x-ray, **miniature photoprints** should be supplied. Photographs should be supplied in high quality glossy paper not larger than 203 mm x 254 mm (8" x 10"). In case of microphotograph, **stains used and magnification should be mentioned**. Each illustration should bear on its back the figure number and an arrow indicating the top. All illustrations should be **colour** and should be submitted in **triplicate** with suitable legends.

**References—**References should be **numbered in the order in which they are first mentioned in the text**. The full list of references at the end of the communication should be arranged in the order mentioned below (names and initials of all authors and/or editors up to 6; if more than 6, list the first 6 followed by *et al.*):

**Reference from journal:**

Cogo A, Lensing AWA, Koopman MMW, Picovella T, Sivagusa S, Wells PS, *et al.*—Compression ultrasonography for diagnostic management of patients with clinically suspected deep vein thrombosis: prospective cohort study. *BMJ* 1998; **316**: 17-20.

**Reference from book:**

Hundin RI—Bleeding and thrombosis. In: Wilson JD, Braunwald E, Isselbacher KJ, Petersdorf RG, Martin JB, Fauci AS, *et al.* editors—Harrison's Principles of Internal Medicine. Vol 1. 12th ed. New York: McGraw Hill Inc, 1991: 348-53.

**Reference from electronic media:**

National Statistics Online—Trends in suicide by method in England and Wales, 1979-2001. [www.statistics.gov.uk/downloads/theme\\_health/HSQ\\_20.pdf](http://www.statistics.gov.uk/downloads/theme_health/HSQ_20.pdf) (accessed Jan 24, 2005): 7-18.

**Only verified references against the original documents** should be cited. Authors are responsible for the accuracy and completeness of their references and for correct text citation. **The number of reference should be kept limited to 20 in case of major communications and 10 for short communications.**

**Dual publication:** If material in a submitted article has been published previously or is to appear in part or whole in another publication, the Editor must be informed.

**Forwarding letter:** The covering letter accompanying the article should contain the name, complete postal address along with Mobile number & E-mail identity of one author as correspondent and must be signed by all authors. The correspondent author should notify change of address, if any, in time.

**Declaration:** A declaration should be submitted stating that the manuscript represents valid work and that neither this manuscript nor one with substantially similar content under the present authorship has been published or is being considered for publication elsewhere and the authorship of this article will not be contested by any one whose name(s) is/are not listed here, and that **the order of authorship as placed in the manuscript is final and accepted by the co-authors**. Declarations should be signed by all the authors in the order in which they are mentioned in the original manuscript.

Matters appearing in the Journal are covered by copyright but no objection will be made to their reproduction provided permission is obtained from the Editor prior to publication and due acknowledgment of the source is made.

— Hony Editor

*Journal of the Indian Medical Association (JIMA),*

*Sir Nilratan Sircar IMA House,*

*53, Sir Nilratan Sircar Sarani (Creek Row), Kolkata-700014*

*Phone: (033) 2236-0573, 2237-8092,*


*E-mail: jmaa1930@rediffmail.com*

*Website: <http://www.ejma.in> & <https://onlinejima.com>*

<sup>1</sup>International Committee of Medical Journal Editors—Uniform Requirements for Manuscripts Submitted to Biomedical Journals. *JAMA* 1997; **277**: 927-34

# Two Feathers in the Cap of Team JIMA in 2019

## JIMA goes SMART




### For JIO Users

- 👉 Download 'JIOCHAT' App
- 👉 Search on JioChannel for 'Journal of IMA'
- 👉 Touch the link you received
- 👉 Download the 'jionews' App
- 👉 Search for 'JIMA' in jionews

### For Non JIO Users

- 👉 Download the "jionews" App
- 👉 Search for 'JIMA' in jionews

INDEX  COPERNICUS  
INTERNATIONAL





Journal title:	
Journal of the Indian Medical Association	
ISSN:	
0019-5847	
GICID:	
n/d	
Country / Language:	
IN / EN	
Publisher:	
Evangel Publishing	
Citation:	14
ICV 2018:	69.74
MNISW 2019:	N/D
ICV 2017:	N/A

Please log on:

<https://journals.indexcopernicus.com/search/details?id=37323&lang=pl>

## Glimpses of NATCON 2019, Kolkata



**JOURNAL OF THE INDIAN MEDICAL ASSOCIATION:**

Sir Nilratan Sircar IMA House, 53 Sir Nilratan Sarkar Sarani (Creek Row), Kolkata - 700 014  
Phones: (033) 2236-0573, 2237-8092, Fax: (033) 2236-6437, E-mail : jima1930@rediffmail.com,  
Website: <http://www.ejima.in>, <https://onlinejima.com> ; [www.ima-india.org](http://www.ima-india.org), [www.ima-india.org/ejima](http://www.ima-india.org/ejima)  
Head office: Indian Medical Association, IMA House, Indraprastha Marg, New Delhi - 110 002  
Telephones: +91-11-2337 0009, 2337 8680, Fax: +91-11-2337 9470, 2337 0375,  
Telegram: INMEDICI, New Delhi - 110 002, Email: [hsg@ima-india.org](mailto:hsg@ima-india.org); Website: [www.ima-india.org](http://www.ima-india.org)

Date of Publication : 15th January, 2020

Registration No. KOL RMS / 476 / 2017-19

RNI Regd. No. 2557/1957  
VOL. 64, NO. 01, January 2020, Kolkata



WHO GMP  
CERTIFIED

[www.jmitra.co.in](http://www.jmitra.co.in)

Setting **GOLD STANDARD** in  
**HIV RAPID** Diagnosis.

4th Generation

**HIV TRI-DOT + Ag**

Rapid Visual Test for Detection of HIV-1 p24 Antigen and  
Differential detection of Antibodies to HIV-1 & HIV-2

First Company  
in India to be granted  
Drug Manufacturing  
Licence for HIV  
Antigen Rapid Test.



p24 Antigen Detection

100%\* Sensitivity

100%\* Specificity

Unique Washing Step

Approved by CDSCO\*\*

\* Evaluated By: National Institute of Biologicals

\*\* Source: [http://cdsco.nic.in/Medical\\_div/List\\_of\\_critical\\_Diagnostic\\_Kits\\_Approved\\_For\\_Blood\\_Bank\\_Use\\_Till\\_Feb\\_200.pdf](http://cdsco.nic.in/Medical_div/List_of_critical_Diagnostic_Kits_Approved_For_Blood_Bank_Use_Till_Feb_200.pdf)

Convenient Packsize: 10 Tests, 50 Tests



Since 1969

**J. Mitra & Co. Pvt. Ltd.**

.....a vision to serve mankind<sup>®</sup>

• Rapid Test Kits • Elisa Test Kits • Confirmatory Tests • Blood Grouping Sera • Fluorescence Immunoassay Test Kits

E-mail: [jmitra@jmitra.co.in](mailto:jmitra@jmitra.co.in) | Tel.: +91-11-471-30-300 | [www.jmitra.co.in](http://www.jmitra.co.in)

If not delivered please return to  
Journal of the IMA (JIMA)  
Sir Nilratan Sircar IMA House,  
53 Sir Nilratan Sarkar Sarani  
(Creek Row), Kolkata - 700 014

Printed and Published by Dr Sanjoy Banerjee on behalf of Indian Medical Association and printed at Prabaha,  
45, Raja Rammohan Sarani, Kolkata 700009, and Published from Sir Nilratan Sircar IMA House, 53 Sir Nilratan  
Sarkar Sarani (Creek Row), Kolkata 700014. Editor : Dr Golokbihari Maji